

T-Z 2.5% w/v Oral Solution

T-Z 2.5% w/v Oral Solution appears to be colourless to light yellow solution packed in a 1 liter white plastic bottle, aluminium sealed cap and label.

Composition:

Each mL contains
Toltrazuril

25 mg

Pharmacodynamics:

Toltrazuril acts as an anticoccidial which is active against all intracellular stages. It is a triazinon derivative. It acts against coccidia of the genus Eimeria. It is active against all intracellular development stages of coccidia of the merogony (asexual multiplication) and gamogony (sexual phase). All stages are destroyed, and hence the mode of action is coccidiocidal.

Pharmacokinetics:

In poultry toltrazuril is absorbed at a rate of at least 50%. The highest equivalent concentrations are found in liver. The active substance is rapidly metabolised in poultry. The main metabolite is characterised as a sulfone-derivative. About 1 week after the last dose this metabolite represents by far the most relevant residue in the animal.

Indications:

For the treatment of Coccidiosis in Broilers and Broiler Breeders. T-Z 2.5% w/v is effective against Eimeria acervulina, E. brunetti, E. maxima, E. necatrix, E. tenella, E. mitis.

Directions for Use:

Add 1 ml of T-Z 2.5% w/v into 1 litre of drinking water (25 ppm) for continuous medication for 48 hours OR add 3 ml of T-Z 2.5% w/v into 1 litre of drinking water (75 ppm) treating at 8 hours per day, on 2 consecutive days or as directed by veterinarian.

Mode of Administration:

Oral administration.

Contraindication:

None.

Statement on Usage During Pregnancy and Lactation:

None to be known of.

Interactions with Other Medicaments:

None known. T-Z 2.5% w/v oral solution is compatible with coccidiostats and amoebostats.

Adverse Effects / Undesirable Effects:

None to be known of.

Overdose and Treatment:

A threefold overdose is well tolerated by calves without signs of intolerance. If the recommended dose is exceeded beyond 3 - 5 times there is a decrease in water intake.

Warning and Precautions:

T-Z 2.5% w/v oral solution does not interfere with the development of immunity against coccidiosis. As with all anticoccidials, prolonged use may result in the development of resistant strains.

Special precautions for use in animals

Dilutions more concentrated than 1:1,000 (1 ml T-Z to 1 litre drinking water) may result in precipitation. Predilution is not recommended.

Special precautions to be taken by the person administering the medicinal product to animals

T-Z is an alkaline solution.

Wear synthetic rubber gloves when handling the product.

Wash any splashes from skin or eyes immediately with water.

Wash hands and exposed skin after use.

Do not eat, drink or smoke whilst using the product.

Other precautions

Only make up sufficient drinking water for requirements.

Wash out empty container with water 3 times when medicating birds and add washings to drinking water.

Store litter from treated houses as long as possible. However, it should only be stored where there is no risk of polluting surface water or groundwater.

Do not apply to agricultural land at a rate greater than 1.5 tonnes/hectare/year.

The principle metabolite of toltrazuril is persistent and mobile in groundwater.

Withdrawal Period (s):

Meat: 18 days

Do not use in birds producing eggs for human consumption.

Storage Condition:

Store below 30°C. Protect from sunlight and heat exposure.

Keep medicine out of reach of children

(Jauhkan ubat dari jangkauan kanak-kanak).

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Shelf life after opened: 28 days

Shelf life after reconstitution: 1 day

Shelf life for sale: 3 years

Packaging Available:

1L.

Date of revision:

23-6-2023

Name and Address of Manufacturer:

NAM PHARMA SDN. BHD.

5, Lebuhr Perusahaan Klebang 11,

Taman Perindustrian Antarabangsa IGB,

31200 Chemor, Perak, Malaysia.

Tel: +605-2931838, 1839 Fax: +605-2931833