

Sulfa-Q 25%w/w Water Soluble Powder

FOR ANIMAL USE ONLY

Product Name :

Sulfa-Q 25%w/w Water Soluble Powder

Name and Strength of Active Substance(s)

Each gram of Sulfa-Q 25%w/w Water Soluble Powder contains Sulfaquinoxaline Sodium 250 mg.

Product Description

Sulfa-Q 25%w/w Water Soluble Powder is a off-white to light yellowish water soluble powder packed in 500g or 1000g aluminum sachets.

Pharmacology

Sulfa-Q 25%w/w Water Soluble Powder contains the sulfaquinoxaline sodium, which is a sulfonamides antimicrobial agent. Mechanism of action : Sulfonamides are structural antagonists of para-aminobenzoic acid (PABA). Sulfonamides interfere with the biosynthesis of folic acid in bacterial cells by competitively preventing PABA from incorporation into the folic acid (pteroylglutamic) acid molecule

Pharmacokinetic

Sulfonamides are well absorbed and well distributed in the body. Levels of drugs tends to be highest in the liver, kidney and lung, and lower in muscle and bone. Sulfonamides cross the placenta and also distributed into milk. Sulfonamides are both renally excreted and metabolized. Renal excretion of unchanged drug occurs via both tubular secretion and glomerular filtration. Mechanisms of metabolism are usually acetylation and glucuronidation. The acetylated metabolites may be less soluble and crystallization in the urine can occur particularly at lower pH.

Indications

For the control of coccidiosis outbreaks in :

- chicken caused by Eimeria tenella, E. necatrix, E. acervulina, E. maxima, and E. brunetti.
- turkey caused by Eimeria meleagris and E. adenoides.

Recommended Dosage

For chickens:

Mix 500 g Sulfa-Q 25%w/w Water Soluble Powder per 300 L of water, treat for 2 days, skip for 3 days, then give 500 g per 500 L of water and treat for 2 more days. Repeat treatment if necessary for 2 more days at 500 g per 500 L of water.

For turkeys:

Mix 500 g Sulfa-Q 25%w/w Water Soluble Powder per 500 L of water and treat for 2 days, skip 3 days, give for 2 days, skip 3 days and give for 2 more days. Repeat treatment if necessary.

Direction of Use

Mixing Directions : To assure thorough dissolution, place the Sulfa-Q 25%w/w Water Soluble Powder in a mixing container and add two gallons of water . Stir until all the powder completely dissolved. Mix this concentrated stock solution with water to make the desired final volume of treated animal drinking water.

Contraindications

Not to be used in patient hypersensitive to sulfaquinoxaline.

Warnings and Precautions

Do not change litter unless absolutely necessary. Do not give flushing mash. Consult a veterinarian or poultry pathologist for diagnosis. May cause toxic reactions unless the drug is evenly mixed in water at dosages indicated and used according to directions. For control of outbreaks of disease, medication should be initiated as soon as the diagnosis is determined. Medicated chickens must actually consume enough medicated water which provides a recommended dosage of approximately 10 to 45 milligrams per pound per day depending on the age, class of animal, ambient temperature, and other factors. Do not give to chickens within 10 days of slaughter for food. Do not medicate chickens producing eggs for human consumption. Make fresh drinking water daily.

Drug Interactions

Combination of sulfaquinoxaline with amprolium or diaminopyrimidines (such as pyrimethamine or trimethoprim) will provide synergistic antibacterial action and enhance spectrum of activity.

Pregnancy and lactation

Do not use in laying birds producing eggs for human consumption.

Adverse Effects

Prolonged administration of sulfaquinoxaline at higher doses may result in depressed feed or water intake, deposition of sulfaquinoxaline crystals in kidney and interference with normal blood clotting.

Overdose and Treatment

Toxicosis from sulfaquinoxaline use in animals is rare. Sulfaquinoxaline has caused hypothermia in animals. These animals responded to vitamin K treatment.

Withdrawal Period

10 days.

Shelf-life (in original packing)

3 years from the date of manufacture. Any opened sachet or unused medicated water to be discarded after 24 hours.

Shelf-life (after opening)

Aluminium sachet is designed for 1 time usage. All powder must be dissolved in drinking water upon opening.

Shelf-life (after opening and reconstitution)

Medicated water must be prepared fresh daily and use within 24 hours. Any unused medicated water is to be discarded after 24 hours.

Maximum Residual Limit (MRL)

Sulfaquinoxaline : Edible offal, muscle (chicken) = 100µg/kg [Ref. REGOV, Version 2, dated Dec 2009]

Storage Conditions

Store in a dry place below 25°C. Keep out of reach of children. Jauhkan dari kanak-kanak.

Disposal of Containers

Traces of medicated powder left in an opened aluminium sachet can be rinsed out with clean animal drinking water and use in animal treatment. Cleaned aluminium sachet can be safely disposed in trash according to local government regulations.

Dosage Forms and packaging available

Available in 500g aluminium sachets packed in 10, 20 or 50 sachets per box and 1000g aluminium sachets packed in 5, 10, 20 or 25 sachets per box.

Name and Address of manufacturer

Life Biopharma Sdn. Bhd. (Company No. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2011-11-23