

Product Name:

TYLOVET 50% Water Soluble powder

Composition:

Each gram contains:

Tyrosin (as tartrate) 500mg

Product Description:

Light Yellow powder

Pharmacodynamics:

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms.

The tylosin spectrum of activity includes Gram-positive bacteria including *Clostridium perfringens* and some Gram-negative strains such as *Pasteurella* and *Mycoplasma* spp. at concentrations of 16g/ml or less.

Pharmacokinetics:

Absorption: Tylosin reaches maximal blood levels between 1 and 3 hours after an oral dose. Minimal or no blood levels remain 24 hours after an oral dose. Distribution: After oral doses were given to pigs, tylosin was found in all tissues, between 30 minutes and two hours after administration, except for the brain and spinal cord. Biotransformation and Elimination: It has been shown that most of the material which is excreted is to be found in the faeces and consists of tylosin (factor A), relomycin (factor D) and dihydrodesmycosin..

Indications:

For the control of *Mycoplasma synoviae* airsacculitis in chickens and *Mycoplasma gallisepticum* in chickens and turkeys. In the field tylosin has also proved useful in reducing the level of infection following stress associated with live vaccination. As an aid in the control of outbreaks of necrotic enteritis in chickens caused by *Clostridium perfringens*. For the prevention and control of enzootic pneumonia, swine dysentery and other scourings caused by organisms (e.g. *Lawsonia intracellularis*) sensitive to tylosin, in pigs. For the control of pneumonia in cattle associated with mycoplasmas and *Pasteurella multocida* sensitive to tylosin.

Dosage and Administration:

For oral administration. Recommendations for use in chickens and turkeys: Tylovet 50% Water Soluble Powder is administered in the drinking water at a concentration of 1 g per litre. As an aid in the control of outbreaks of necrotic enteritis caused by *Clostridium perfringens* in chickens use Tylovet 50% Water Soluble Powder in the drinking water for 5 days at a concentration of 0.3g per litre water, to provide 20-50 mg/kg bw, depending on the age and the water consumption of the birds.

Recommendations for use in pigs:

1. Tylovet 50% Water Soluble Powder is administered in the drinking water to provide 25 mg tylosin/kg bodyweight. This may normally be achieved by adding 0.5g per litre.
2. A medicated solution of drinking water should generally be administered until 24 hours after scouring or respiratory symptoms have ceased, normally 3-10 days. The diagnosis should be reviewed if there is no response after 5 days of medication.

Recommendations for use in calves:

1. One gram of tylosin activity per calf administered orally twice daily for seven to fourteen days. The tylosin tartrate may be incorporated into the milk or reconstituted milk replacer at the time of feeding.
2. One gram of tylosin activity should be incorporated in milk or milk replacer twice daily for each calf. This should be continued for 7-14 days dependent on response. The intake of medicated water depends on the clinical condition of the animals. In order to obtain the correct dosage, the concentration of Tylovet 50% WSP has to be adjusted accordingly.

Contraindications:

Not to be used in animals with known sensitivity to the active ingredient.

Do not leave or dispose of water containing tylosin tartrate where it may be accessible to either animals not under treatment or to wildlife

Interaction with Other Medicaments:

None known.

Pregnancy and Lactation:

No adverse effects to tylosin have been seen in fertility, multi-generation or teratology studies.

Side effects:

None known.

Symptoms and Treatment of Overdose:

There is no evidence of tylosin toxicity in animals, at dose rates of up to 1000 mg/kg.

Warning and Precautions:

Keep out of reach of children.

Jauhi dari kanak-kanak.

Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided. To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator or a non-disposable respirator. Wash hands after use. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water. Do not handle the product if you are allergic to ingredients in the product. If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Withdrawal Periods:

Meat & offal

Pig:	7days
Broller and pullets:	7days
Calves:	14 days

Do not use in laying hens producing eggs for human consumption.

Storage conditions:

Store in cool, dark place. Store below 30° C. Protect from direct light.

Pack Size:

1kg, 100gm

Distributed by Product Registration Holder:

LIM SENG PHARMACHEM SDN. BHD.

2692, Tingkat Perusahaan Enam,

Prai Industrial Estate MK1,

13600 Prai, Pulau Pinang, Malaysia.

Manufactured By:

NAM PHARMA SDN. BHD.

5, Lebuhr Perusahaan Klebang II,

Taman Perindustrian Antarabangsa IGB

31200 Chemor, Perak.

DATE OF REVISION: APRIL 2019