

SPECTRO 5% W/V ORAL SOLUTION

SPECTRO 5% W/V ORAL SOLUTION appears to be a red solution.

Composition

Each mL contains Spectinomycin 50 mg
(as *Spectinomycin hydrochloride*)

Pharmacodynamics

Spectinomycin is an aminocyclitol antimicrobial and has been shown to be produced by two kinds of fungi, *Streptomyces flavopersicus* and *Streptomyces spectabilis*. It is similar to the aminoglycosides, acting by binding to ribosomal 30s subunits of the bacteria. In-vitro tests have shown spectinomycin to be effective against a range of Gram-negative and Gram-positive bacteria, although in the field its spectrum of action is for Gram-negative bacteria. Anaerobic organisms are generally resistant. It has been demonstrated that spectinomycin is more bacteriostatic than bactericidal. The mechanism of action appears to be by inhibition of protein synthesis.

Pharmacokinetics

Spectinomycin is poorly absorbed from the alimentary canal. In the blood, protein binding is low at less than 10%. It is almost totally eliminated by renal glomerular filtration with 75% or more cleared in 4 hours.

Indications

For the treatment and control of enteritis caused by strains of *Escherichia coli* sensitive to spectinomycin in piglets.

Dosage and Administration

By oral administration.

Piglets under 4.5 kg (10 lb): 1 pump (1 ml) twice daily.
Piglets between 4.5 kg (10 lb) and 7.0 kg (15 lb): 2 pumps (2 ml) twice daily.

Treatment may be continued twice daily for 3-5 days.

If pigs do not improve within 48 hours, re-diagnosis is suggested

Insert the doser in the bottle and push the piece of clear plastic tubing over the end of the pump spout. Press the plunger a few times to fill both the pump and plastic extension tube with medication.

The preparation should be administered well back on the tongue. Each pump of the plunger delivers 1 ml of solution containing 50 mg of spectinomycin. When not in use, remove the dosing pump from the bottle. Place the clear plastic tube in the neck of the bottle and press the plunger a few times to return any medication in the pump to the bottle. Before storing for future use, always rinse out the doser with water to prevent the pump parts from sticking.

Contraindications

Known hypersensitivity to the active ingredient.

Do not use in pigs over 4 weeks of age or over 7 kg bodyweight.

Use with caution in animals suffering from renal damage.

Special warnings for each target species

None.

Special precautions for use

Special precautions for use in animals

Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local (regional, farm level) epidemiological information about susceptibility of the target bacteria.

Special precautions to be taken by the person administering the medicinal product to animals

Do not handle in case of known hypersensitivity to spectinomycin. In case of accidental splashing on to the skin or eyes, wash the affected area with plenty of clean running water. Wash hands thoroughly after use.

Interactions with Other Medicaments

None known.

Pregnancy and Lactation

Not applicable.

Side Effects

None known.

Overdose and Treatment

Because the product is so poorly absorbed from the alimentary canal, it is difficult to overdose.

The symptoms of overdosage in piglets are not known. There is no known antidote to overdosage.

Withdrawal Period(s)

Animals must not be slaughtered for human consumption during treatment.

Pigs may be slaughtered for human consumption only after 12 days from the last treatment.

Storage

Store below 30°C. Protect from heat and sunlight exposure.

Keep out of reach of children/ Jauhkan daripada anak-kanak.

Any unused product or packaging should be disposed of in accordance with local legislation.

Shelf life packaged for sale: 3 years.

Shelf life after first opening: 30 days.

Packaging Available

100ml

Manufacturer and Product Holder

NAM PHARMA SDN. BHD.

5, Lebuhr Perusahaan Klebang 11,
Taman Perindustrian Antarabangsa IGB,
31200 Chemor, Perak, Malaysia.

Date of revision

02.03.2026