

HEMOFORT 20% INJECTION

Description and Composition

HEMOFORT 20% INJECTION is a dark brown injection solution. Each ml contains 200mg Iron (as dextran) as its active ingredient.

Pharmacodynamics

Iron (as dextran) is an anti-anemic acting significantly increasing the hemoglobin. After administration, the iron is stored in the tissues, and then gradually released from the complex with dextran. It is incorporated in the form of hemoglobin proteins such as ferritin and hemosiderin. Iron is administered as an iron dextran complex to avoid the toxic effects of free iron.

Pharmacokinetics

After administration, about 60% of iron is absorbed after 3 days and 90% after 1 to 3 weeks.

Indication

For the prevention and treatment of iron deficiency anemia in swine.

Recommended Dose

200 mg of iron per piglet by deep intramuscular injection preferably in the thigh muscles in the three days following the birth or 1 ml at one injection.

Route of Administration

By intramuscular injection.

Contraindication

- Do not use in animals with known hypersensitivity to the active ingredient.
- Do not use in older pigs as ham staining may occur in animals over 4 weeks of age.

Warning and Precaution

Special warnings for each target species.

No

Special precautions for use

i) Special precautions for use in animals

Strictly aseptic technique during injection.

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals.

In case of accidental self - injection, seek medical advice immediately and show the label.

iii) Other precautions.

No.

Interaction with Other Medicinal Product

None known

Pregnancy and Lactation

NA

Side Effect

Staining of injection.

Occasionally , cases of anaphylactic shock , some fatal, have been reported following administration of iron (as dextran)

Symptoms and Treatment of Overdose

Not applicable provided the recommended dosage is not exceeded.

Incompatibilities

None known

Storage Condition

Store below 30°C. Protect from light.

Shelf Life

3 years. Following withdrawal of the first dose, use the product within 24hours. Any unused material should be discarded.

Withdrawal Period

Meat and offal: Zero days

Packing

100ml

REVISED: 12.6.14

	Manufactured by:
	Range Pharma Sdn Bhd 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: sales@rangepharma.com.my