

USER INFORMATION

InfuVet HM

COMPOUND SODIUM LACTATE (HARTMANN'S SOLUTION) INTRAVENOUS INFUSION BP FOR VETERINARY USE

Composition:

The Solution contains:		Electrolytes :	mmol/L
Sodium Chloride BP	0.600% w/v	Sodium Ion (Na ⁺)	131.0
Potassium Chloride BP	0.040% w/v	Potassium Ion (K ⁺)	5.0
Calcium Chloride Dihydrate BP	0.027% w/v	Calcium Ion (Ca ⁺⁺)	2.0
Sodium Lactate USP	0.320% w/v	Chloride Ion (Cl ⁻)	111.0
		Lactate Ion (HCO ₃ ⁻)	29.0
		Osmolarity :	278 mOsm/L

Characteristics:

InfuVet HM (Compound Sodium Lactate (Hartmann's Solution) Intravenous Infusion BP) is a colourless solution.

Pharmacodynamics:

The veterinary medicinal product is administered to replace or prevent dehydration, to correct acid-base, fluid and electrolyte abnormalities in different clinical conditions. The electrolytes Na⁺, K⁺, Ca⁺⁺, Cl⁻ as well as the metabolisable anion lactate are indispensable for the maintenance and correction of fluid and electrolyte homeostasis and acid-base balance. All substrates are occurring during normal physiological metabolism.

Pharmacokinetics:

Due to intravenous administration the bioavailability of the active substances is 100%. The metabolism of this veterinary medicinal product corresponds to each of its components: Na⁺, K⁺, Ca⁺⁺, Cl⁻ and lactate. Sodium lactate decomposes into bicarbonate, which later is converted into pyruvate that is used in the Krebs tricarboxylic cycle.

Target Species:

Cats, Dogs, Equine, Swine (Pig), Sheep, Goats, Cattle

Indication:

Indication for all target animal species:

- Isotonic dehydration
- Metabolic acidosis
- Hypotonic dehydration
- Maintenance of normal extracellular fluid levels
- Electrolyte replacement in burns

Recommended Dosage:

Intravenous use.

The volume and rate of infusion will depend upon the clinical condition, existing deficits of the animal, maintenance needs and continuing losses.

Generally aim to correct hypovolaemia by 50% initially (ideally over 6 hours but faster if necessary) and reassess by clinical examination.

Deficits are generally in the range of 50 ml/kg (mild) to 150 ml/kg (severe). An infusion rate of 15 ml/kg/bw/hour is recommended in the absence of shock (range 5-25 ml/kg/bw/hour).

In shock, high initial infusion rates, up to 90 ml/kg/bw/hour, are needed. High infusion rates should not be continued for longer than 1 hour unless urine output is restored. The maximum infusion rate should be decreased in the presence of cardiac, renal and pulmonary disease.

Do not use if container or closure is damaged. For single use only. Solutions containing visible solid particles should not be administered.

Route of administration : Intravenous (IV)

Contraindications:

Do not use in animals with:

- Alkalosis of any origin
- Oedema (hepatic, renal, or cardiac)
- Overhydration
- Hyperkalaemia, hypernatraemia, hyperlactataemia
- Hepatic insufficiency

Warnings and Precautions:

Special precautions for use in animals

Before administering this solution the clinical and biological data of the animal have to be carefully examined.

Monitoring of serum electrolyte levels should be obliged in cases of electrolyte imbalances, such as hypertonic or hypotonic dehydration, or a single increase of one electrolyte (e.g. hyperchloraemia) as well. Furthermore monitoring of the acid-base and the clinical condition of the animal should accompany the treatment with this veterinary medicinal product.

During use of this veterinary medicinal product, the fluid volume range must be considered. Infusion of larger than necessary volumes may lead to cardiovascular overload and pulmonary oedema.

This veterinary medicinal product should be used with caution in congestive heart failure, severe renal insufficiency and in animals treated with corticoids and derivatives.

Due to the potassium content of this solution it should be used prudently in severe renal impairment.

Infusion of this solution containing lactate ions may cause metabolic alkalosis.

In animals with liver function disorders, the solution may cause acidosis because degradation of lactate into bicarbonate requires an intact liver metabolism.

Slow infusion into a large blood vessel should be performed under conditions of strict asepsis. Do not inject intramuscularly.

During treatment clinical and biological state of the animal should be monitored.

The solution should be administered at body temperature. Warm up the solution only by immersion in hot water (< 40°C).

Caution:

Single dose container. Discard all unused contents or when the seal is first broken. Do not use if the solution is discoloured and leakage is detected. If any visible solid particles, growth or turbidity appears during storage, the product should not be used. Before use check for minute leaks by squeezing bag firmly.

Interactions with Other Medicaments:

The compatibility of any additive should be checked before use.

Statement on Usage During Pregnancy and Lactation:

The safety of the veterinary medicinal product has not been established during pregnancy or lactation. Use only accordingly to the benefit/risk assessment by the responsible veterinarian.

Adverse Effects/Undesirable Effects:

This veterinary medicinal product contains calcium, thus an effect on the heart cannot be ruled out.

Overdose and Treatment:

Overdose may result in cardiovascular overload and pulmonary oedema, which can lead to following symptoms such as restlessness, coughing and polyuria. In case overdose has occurred the rate of infusion should be drastically reduced or the infusion should be stopped.

Withdrawal Period(s):

Meat and offal: Zero days

Milk: Zero hours

Storage Condition:

Do not store above 30°C.

Direction for use:

For use only by or under the supervision of a veterinary surgeon.

Incompatibilities:

This veterinary medicinal product is incompatible with Chlortetracycline, Amphotericin B and Oxytetracycline.

Mixtures with additives and other drugs (e.g. oxalate-, phosphate- and carbonate- / hydrogen carbonate- containing ones) may cause incompatibilities.

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

Shelf life:

3 years from manufacturing date in the proposed storage condition in Medical Collapsible Bag.

3 years from manufacturing date in the proposed storage condition in LDPE Bottle.

Do not use after expiry.

Dosage form and packaging available:

- 1000mL X 10 LDPE bottles per box
- 1000mL X 10 medical collapsible bags per box
- 3000mL X 4 medical collapsible bags per box
- 5000mL X 1 medical collapsible bag per box.

Disposal of containers:

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Manufacturer/ Product Registration Holder:



AIN MEDICARE SDN.BHD
Jalan 6/44, Kaw. Perindustrian Pengkalan Chepa 2, 16100 Kota Bharu, Kelantan, MALAYSIA

Date of Revision : 24.06.2014