

USER INFORMATION
InfuVet NS
SODIUM CHLORIDE 0.9% w/v INTRAVENOUS INFUSION BP
FOR VETERINARY USE

Composition:
Sodium Chloride BP 9 g/L

Electrolytes (approx):	mmol/L	(mEq/L)
Sodium Ion (Na ⁺)	150	150
Chloride Ion (Cl ⁻)	150	150
Osmolarity :	308 mOsm/L	

Characteristics:
Sodium Chloride 0.9% Intravenous Infusion BP is an isotonic solution containing 150 mmol each of Sodium and Chloride per litre. The solution is clear and colourless.

Pharmacodynamics:
The intravenous infusion is used to replace depleted water and electrolytes when fluids cannot be given orally. It will restore extracellular volume (including plasma volume) and, where metabolic alkalosis is present, it will help to correct it.

Pharmacokinetics:
Intravenous infusion ensures rapid distribution. The constituents of the infusion solution will be metabolised and excreted through the same pathways as those substances derived from normal dietary sources.

Incompatibilities:
None known

Target Species:
Cats, Dogs, Equine, Cattle

Indication:
This product is administered by intravenous infusion for the treatment of dehydration in cats, dogs, equine, cattle. It may be used to correct hypovolaemia resulting from shock or gastrointestinal disease (especially where metabolic alkalosis is present, e.g. in cases of sustained vomiting or abomasal disorders in cattle). It may be administered to meet normal fluid and electrolyte requirements when fluids cannot be given orally

Recommended Dosage:
Intravenous use.
The volume and rate of infusion will depend upon the clinical condition, existing deficits of the animal, maintenance needs and continuing losses.
Generally aim to correct hypovolaemia by 50 % initially (ideally over 6 hours but faster if necessary) and reassess by clinical examination.
Deficits are generally in the range of 50 ml/kg (mild) to 150 ml/kg (severe). An infusion rate of 15 ml/kg/hour is recommended in the absence of shock (range 5-75 ml/kg/hour).
In shock, high initial infusion rates, up to 90 ml/kg/hour, are needed. High infusion rates should not be continued for longer than 1 hour unless urine output is restored. The maximum infusion rate should be decreased in the presence of cardiac, renal and pulmonary disease.

Route of administration : Intravenous (IV)

Contraindications:
None

Warnings and Precautions :
Special precautions for use in animals:
Do not use unless the solution is clear, free from visible particles and the container is undamaged.
Sodium overload may occur in animals with cardiac or renal impairment. It should be noted that sodium excretion may be impaired post-surgery/trauma.
A risk of thrombosis with intravenous infusion should be considered.
The product should be warmed to approximately 37°C prior to the administration of large volumes, or if the administration rate is high, in order to avoid hypothermia.
This product should not be used for longer than is necessary to correct and sustain circulating volume. Inappropriate / excessive use may worsen or create a metabolic acidosis.
Maintain aseptic precautions.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:
None.

Caution:
Single dose container. Discard all unused contents or when the seal is first broken. Do not use if the solution is discoloured and leakage is detected. If any visible solid particles, growth or turbidity appears during storage, the product should not be used. Before use check for minute leaks by squeezing bag firmly.

Interactions with Other Medicaments:
Compatibility of the solution should be checked before used

Statement on Usage During Pregnancy and Lactation:
Use under veterinary supervision

Adverse Effects/Undesirable Effects:
Excessive infusion rates can cause restlessness, moist lung sounds, tachycardia, tachypnoea, nasal discharge, coughing, vomiting and diarrhoea.

Overdose and Treatment:
Monitor fluid output. The administration of a diuretic may be necessary.

Withdrawal Period(s):
Cattle and horses:
Meat and offal: Zero days
Milk: Zero hours

Storage Condition:
Do not store above 30°C.

Direction for use:
For use only by or under the supervision of a veterinary surgeon.

Shelf life:
3 years from manufacturing date in the proposed storage condition in Medical Collapsible Bag.
5 years from manufacturing date in the proposed storage condition in LDPE Bottle.
Do not use after expiry.

Dosage form and packaging available:
1000mL X 10 LDPE bottles per box
1000mL X 10 medical collapsible bags per box
3000mL X 4 medical collapsible bags per box
5000mL X 1 medical collapsible bag per box.

Disposal of containers:
Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Manufacturer/ Product Registration Holder:
 AIN MEDICARE SDN.BHD.
Jalan 6/44, Kaw, Perindustrian Pengkalan Chepa 2, 16100 Kota Bharu, Kelantan, MALAYSIA.

Date of Revision : 24.06.2014