

For Veterinary Use Only

Lincomix™ 110
PREMIX

5 kg

Keep out of reach of children
Jauhkan ubat dari kanak-kanak
Controlled Medicine / Ubat Terkawal

Each kg contains:
Lincomycin (as hydrochloride) ----- 110 g

Product Description: A grayish white to light brown powder which has a cereal odor.

Pharmacodynamics: Lincomycin is a lincosamide antibiotic and is active against a wide range of Gram positive aerobic and anaerobic bacteria, anaerobic Gram-negative bacteria and mycoplasma. Lincosamides are generally bacteriostatic. The mode of action is inhibition of protein synthesis at the ribosomal 50S sub-unit level. It is active against the primary causal agent of necrotic enteritis, *Clostridium perfringens*. It is active against the primary causal agent of swine dysentery, *Serpulinae hyodysenteriae* and the exacerbating anaerobes, *Bacteroides*, *Fusobacterium* and *Clostridium spp.*, and also of mycoplasma (enzootic) pneumonia in pigs, *Mycoplasma hyopneumoniae*.

Pharmacokinetics: Levels of Lincomycin in the small intestine of chickens administered 4.4 g lincomycin per 1000 kg feed should approximate to the minimum inhibitory concentrations of sensitive *C. perfringens*. Lincomycin has approximately 50% systemic bioavailability from oral administration in pigs and is significantly re-circulated via biliary route.

Indication:

Swine:

- To reduce the severity of mycoplasma pneumonia in growing swine.
- For the treatment of swine dysentery (bloody scours) associated with *Brachyspira* (*Serpulina*) *hyodysenteriae* in growing swine and for the control of the disease following treatment.
- As an aid in reducing the clinical severity of Porcine Proliferative Enteropathy (Ileitis) associated with *Lawsonia intracellularis*.

Chicken (broilers):

- For control of necrotic enteritis caused by *Clostridium* species or other susceptible organisms.

Recommended Dose:

Swine:

To reduce the severity of *Mycoplasma pneumoniae* in growing swine.

Directions for Use: Complete feeds containing lincomycin base at 220 grams per tonne of complete feed, should be fed to growing swine as the sole ration for 21 days.

Mixing Directions: Add 2 kg of LINCOMIX 110 to each metric tonne of complete feed. The resultant feed will contain 220 g lincomycin base per tonne.

For the treatment of Swine Dysentery (Bloody Scours) associated with *Brachyspira* (*Serpulina*) *hyodysenteriae* in growing swine, and for the control of the disease following treatment.

Directions for Use: Complete feeds containing lincomycin at the following dosage levels should be fed to growing swine as the sole ration. For treatment where pigs have already begun to show signs of disease, provide 110 grams of lincomycin base per tonne of complete feed for 21 days, or until signs of disease (watery, mucoid or bloody stools) disappear, then reduce the medication level to 44 grams of lincomycin base per tonne and continue for a further 4 weeks. Retreat with the above dosage regimen if the symptoms of swine dysentery reappear.

Mixing Directions: Add 1 kg of LINCOMIX 110 to each metric tonne of complete feed. The resultant feed will contain 110 g of lincomycin base per tonne.

Control Level: Add 0.4 kg of LINCOMIX 110 to each metric tonne of complete feed. The resultant feed will contain 44 g of lincomycin base per tonne.

As an aid in reducing the clinical severity of Porcine Proliferative Enteropathy (Ileitis) associated with *Lawsonia intracellularis*.

Directions for Use: Feed 110 grams of lincomycin per tonne of complete feed as the sole ration for 21 days.

Mixing Directions: Add 1 kg of LINCOMIX 110 to each metric tonne of complete feed. The resultant feed will contain 110 g of lincomycin base per tonne.

Chicken (broilers):

For control of necrotic enteritis caused by *Clostridium* species or other susceptible organisms.

Directions for Use: Feed 2 grams of lincomycin per ton of feed

Mixing Directions: Add 18.18 g of LINCOMIX 110 to each metric tonne of complete feed. The resultant feed will contain 2-4 g of lincomycin base per tone.

Route of Administration: To be administered orally via the feed.

Contraindications: Do not administer to animals with known hypersensitivity to the active ingredient. Not for use in animals other than the target species. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feed containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects. Not for use in breeding swine. Do not use in feed containing pellet-binding agents except Lignin Sulfonate.

Warning and Precautions:

To ensure thorough dispersion of the product it should first be mixed with a suitable quantity of feed before incorporation into the final mix.

Persons known to be hypersensitive (allergic) to lincomycin and/or rice husk should not handle this product.

Care should be taken not to inhale any dust. The wearing of approved dust masks (either a disposable half-mask respirator conforming to European Standard EN 149 or a nondisposable respirator to European Standard EN 140 with a filter to EN 143) and safety glasses is recommended during the handling and mixing of this product.

Skin contact should be avoided.

Wash hands and any exposed skin with soap and water immediately after use.

Interaction with other medicaments: Clinical incompatibility may exist between lincomycin and erythromycin due to competitive binding at the ribosomal site of action.

Pregnancy and Lactation: Not applicable.

Side effect: Loosening of faeces and/or mild swelling and irritation of the anus and/or vulva has been occasionally observed within the first two days after onset of treatment. This is usually transient. Very rarely some pigs may show reddening of skin and mild irritable behaviour. These conditions are usually self-correcting within 5-8 days without discontinuing lincomycin treatment.

Overdose and Treatment: Not applicable.

Withdrawal Period: Swine treated at the recommended level of 220 grams of lincomycin per tonne must not be slaughtered for use in food for at least 24 hours after the latest treatment with this drug. No drug withdrawal period before slaughter is required for feed medicated with LINCOMIX 110 g/kg Powder at the recommended level of 44 or 110 grams of lincomycin base per tonne.

Storage Condition: Do not store above 25°C (controlled temperature room). Store in a dry place. Store opened bag in a dry place to prevent caking.

Dosage form and packaging available: Lincomix 110 g/kg Premix is available in 5kg.

Disposal: Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Manufactured by:

Zoetis Suzhou Manufacturing Co., Ltd.

No 690, Jian Lin Road,
Suzhou New District,
Jiangsu 215151, P.R. China.

zoetis™

Product Registration Holder:

Zoetis Malaysia Sdn. Bhd. (8898-V)

Lot 3.1, Level 3, Tower 7,
Avenue 3, Bangsar South,
No. 8, Jalan Kerinchi, 59200,
Kuala Lumpur, Malaysia.

Date of revision: Dec 2018

MAL1407011HA

LOT No.:

MFG Date:

EXP Date: