

|                                                                                                                                     |                                     |
|-------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------|
| <b>VETOQUINOL</b>                                                                                                                   | <b>Nom du produit :</b> Tolfine MY  |
| <b>Créé par :</b> P.JARDEL/M.MARQUES                                                                                                | <b>Référence article :</b> 0521 B L |
| <b>le :</b>                                                                                                                         | <b>Dimensions :</b> 160 x 120 mm    |
| <b>Visa :</b>                                                                                                                       | <b>1 couleur</b>                    |
| <div data-bbox="692 419 768 453" data-label="Image"> </div> <div data-bbox="699 457 764 480" data-label="Text"> <p>281 U</p> </div> |                                     |

The product may be used during lactation. As there are limited data to support use during pregnancy, use of the product in pregnant animals should be at the discretion of the veterinary surgeon after the risks and benefits have been considered.

#### **ADVERSE EFFECTS**

Transient swelling and inflammation may occur at the injection point.

#### **OVERDOSE AND TREATMENT**

None

#### **FURTHER INFORMATION**

Keep out of the reach of children.

Following withdrawal of the first dose, the product should be used within 28 days.

Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

#### **STORAGE CONDITIONS**

Store below 30°C (room temperature)

#### **DOSAGE FORMS AND PACKAGING**

Amber type I glass vials of 50 or 100 ml closed with a type I chlorobutyl rubber stopper.

**REG NUMBER : MAL14070115HA**

**FOR ANIMAL TREATMENT ONLY  
CONTROLLED MEDICINE - UBAT TERKAWAL**

**NAME AND ADDRESS OF MANUFACTURER:  
VETOQUINOL S.A**

Magny-Vernois, 70200 Lure, France

**NAME AND ADDRESS OF REGISTRATION HOLDER:  
GLADRON CHEMICALS SDN. BHD.**

No.7, Jalan TP7, UEP Industrial Park, 40400 Shah Alam,  
Selangor Darul Ehsan, Malaysia

**DATE OF REVISION:** May 2021



**TOLFINE®**

**4% solution for injection  
for cattle and pigs**

#### **COMPOSITION**

Tolfenamic acid..... 4.0% w/v

Benzyl alcohol..... 1.04 % w/v

#### **PRODUCT DESCRIPTION**

Solution for injection, vials of 50ml or 100 ml of clear, colourless to slightly yellow, slightly viscous solution.

#### **PHARMACOLOGICAL PROPERTIES**

**Tolfenamic acid** (N-(2-methyl-3-chlorophenyl) anthranilic acid) is a non-steroidal anti-inflammatory drug (NSAID) belonging to the fenamate group. Tolfenamic acid exerts anti-inflammatory, analgesic and antipyretic activities. The anti-inflammatory activity of tolfenamic acid is mainly due to an inhibition of cyclo-oxygenase and thus to a reduction in the synthesis of prostaglandins and thromboxanes, which are important inflammatory mediators.

#### **PHARMACOKINETIC PROPERTIES**

##### **PIGS**

In pigs, tolfenamic acid injected by IM route at a dose of 2mg/kg is rapidly absorbed from the injection site with a mean maximum plasma concentration of about 2.3µg/ml obtained at about 1 hour. The volume of distribution is approximately 1.3 l/kg in pigs. It is extensively bound to plasma albumin (>97%).

Tolfenamic acid distribution involves extracellular fluids where concentrations similar to plasma are achieved both in healthy and inflamed peripheral tissues. Tolfenamic acid undergoes extensive enterohepatic recirculation and, as a result, prolonged concentrations are found in plasma. The elimination half life varies from 3 - 5 hours in pigs.

## CATTLE

Tolfenamic acid is distributed in all the organs with a high concentration in the plasma, digestive tract, liver, lungs and kidneys. However, the concentration in the brain is low. Tolfenamic acid and its metabolites do not cross the placental barrier to any great extent.

Tolfenamic acid distribution involves extracellular fluids where concentrations similar to plasma are achieved both in healthy and inflamed peripheral tissues. It also appears in milk in the active form, mainly associated with the curds.

Tolfenamic acid undergoes extensive enterohepatic recirculation and, as a result, prolonged concentrations are found in plasma. The elimination half life varies from 8 - 15 hours in cattle.

In cattle and pigs, tolfenamic acid is eliminated mainly unchanged in faeces (~30%) and urine (~70%).

## INDICATIONS, ADMINISTRATION, RECOMMENDED DOSAGE AND WITHDRAWAL

Tolfine is an anti-inflammatory, antipyretic and analgesic agent indicated:

| Species | Indication                                                         | Route | Dose                        | Withdrawal Periods                                       |
|---------|--------------------------------------------------------------------|-------|-----------------------------|----------------------------------------------------------|
| Pig     | Metritis mastitis                                                  | IM*** | 2mg/kg                      | Meat: 3 days                                             |
| Cattle  | Acute mastitis used in conjunction with antibacterial therapy      | IV    | 4mg/kg<br>b w ;<br>1ml/10kg | Meat: 3 days<br>Milk: 24 hours**                         |
|         | Respiratory disease used in conjunction with antibacterial therapy | IV    | 2mg/kg<br>bw;<br>1ml/20kg*  | Meat: 3 days<br>Milk: 24 hours**                         |
|         |                                                                    | SC    | 2mg/kg<br>bw;<br>1ml/20kg*  | Meat: 7 days<br>Milk: <b>not for use in dairy cattle</b> |

*\*may be repeated once after 48 hours. If a second injection is needed this should be given in the opposite side of the animal's body.*

*\*\*milk from treated cattle must only be used for human consumption after 24 hours i.e. at the second milking. Milk from the first milking after treatment should be discarded*

*\*\*\*in neck musculature*

## CONTRA-INDICATIONS

Not for IM administration in cattle

Not for IV or SC administration in pigs

Not for SC administration in dairy cows

## WARNINGS

- Do not exceed 20 ml per intramuscular injection site.
- Do not exceed the stated dose or duration of treatment.
- Use aseptic precautions when administering the product.
- When treating a number of animals the use of a draw-off needle is recommended to avoid excessive broaching of the closure.
- Use is contra-indicated in animals suffering from cardiac, hepatic or renal disease, where there is the possibility of gastro-intestinal ulceration or bleeding, or where there is evidence of hypersensitivity to the product.
- Use in any animal less than 6 weeks of age or in aged animals may involve additional risk. If such use cannot be avoided animals may require a reduced dosage and careful clinical management.
- Avoid use in any dehydrated, hypovolaemic or hypotensive animal as there is a potential risk of increased renal toxicity.
- It is preferable that a NSAID which inhibits prostaglandin synthesis is not administered to animals undergoing general anaesthesia until fully recovered.
- Concurrent administration of potentially nephrotoxic drugs should be avoided.
- Take care to avoid self-injection. In case of eye or skin contact, wash immediately with water.

## INTERACTIONS WITH OTHER MEDICAMENTS

Do not administer other NSAIDs concurrently or within 24 hours of each other. Tolfenamic acid is highly bound to plasma proteins and may compete with other highly bound drugs which can lead to toxic effects.

## PREGNANCY AND LACTATION