

[Draft of Package Insert]

For Animal Use Only

ISOTYL 200

Medicated Premix Powder

CONTENT:

Tilmicosin.....200 g/kg

PRODUCT DESCRIPTION: An almost white powder.

PHARMACODYNAMICS:

Tilmicosin is a semi-synthetic antibiotic of the macrolide group, and is believed to affect bacterial protein synthesis. It is bacteriostatic but at high concentrations it may be bactericidal. Tilmicosin has an *In-vitro* antibacterial spectrum that is predominantly Gram-positive with activity against certain Gram-negative microorganisms. Activity against several mycoplasma species has also been demonstrated.

PHARMACOKINETICS:

Following oral administration, Tilmicosin is distributed throughout the body, but especially high levels are found in the lung and in lung tissue macrophages. It is also distributed in the liver and kidney tissues. Tilmicosin is excreted mainly via the bile into the faeces, but a small proportion is excreted via the urine.

INDICATION:

For the prevention and treatment of pneumonia in swine caused by *Actinobacillus pleuropneumoniae*, *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and other organisms sensitive to Tilmicosin.

DOSAGE & ADMINISTRATION:

The uptake of medicated feed depends on the clinical condition of the animals. In order to obtain a correct dosage, the concentration of tilmicosin has to be adjusted accordingly.

Use the following formula:

$$\text{Kg Premix/tonne feed} = \frac{\text{Dose rate (mg/kg bodyweight)} \times \text{bodyweight (kg)}}{\text{Daily feed intake (kg)} \times \text{premix strength (g/kg)}}$$

Pigs

Administer in the feed at a dose of 8 to 16 mg/kg body weight/day of tilmicosin (equivalent to 200 to 400 ppm in the feed) for a period of 15 to 21 days.

Indication	Dose of tilmicosin	Duration of treatment	Inclusion rate in feed
Prevention and treatment of respiratory disease	8-16 mg/kg bodyweight /day	15-21 days	1-2 kg Isotyl 200/tonne

CONTRAINDICATIONS:

Do not use in any feed (supplement, concentration, or complete feed) containing bentonite. Bentonite in feeds may affect the efficacy of tilmicosin.

Do not use in case of hypersensitivity to tilmicosin or to any of the excipients.

WARNINGS / PRECAUTIONS:

Do not allow horses or other equines access to feeds containing Tilmicosin.

Slurry from treated animals should not be deposited on the same field in successive years.

Operator warnings:

May cause irritation or sensitization by skin contact. Avoid direct skin contact. Wear overalls and impervious gloves when mixing and handling the product. Wash affected parts if skin contact occurs.

Avoid contact with eyes. If accidental eye contact occurs, immediately rinse thoroughly with water.

Cross-resistance between tilmicosin and other macrolide antibiotic has been observed. Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies. Inappropriate use of the product may increase the prevalence of bacteria resistant to tilmicosin and may decrease the effectiveness of treatment with tilmicosin related substances.

DRUGS INTERACTIONS: Do not use simultaneously with other macrolides and lincosamides. Do not mix into feed containing bentonite.

PREGNANCY AND LACTATION: The safety of Tilmicosin in pregnant swine and animals intended for breeding purposes has not been established.

ADVERSE EFFECTS: Occasionally, feed intake may decrease (including feed refusal) in animals receiving medicated feed. This effect is transient.

OVERDOSE AND TREATMENT:

No symptoms of overdose have been seen in pigs fed a ration containing levels of tilmicosin up to 80 mg/kg bodyweight (equivalent to 2000 ppm in the feed or ten times the recommended dose) for 15 days.

STORAGE CONDITIONS: Store below 30°C.

WITHDRAWAL PERIOD(S):

Pigs: 21 days

DOSAGE FORMS AND PACKAGING AVAILABLE: 1kg powder packed into aluminium sachet.

DATE OF REVISION OF PACKAGE INSERT: 09 JULY 2014.

Manufactured by:

PAHANG PHARMACY SDN. BHD. (10826-K)

LOT 5979, JALAN TERATAI,
5½ MILE OFF JALAN MERU, 41050 KLANG,
SELANGOR D.E., MALAYSIA.

Product Registration Holder:

GENPHARMA SDN. BHD. (268066-W)

LOT 5979, JALAN TERATAI,
5½ MILE OFF JALAN MERU, 41050 KLANG,
SELANGOR D.E., MALAYSIA.