

Neomix™ 325

SOLUBLE POWDER

500 g

Package Content: 357.5 g Neomycin sulfate per 500 g product (250 g neomycin activity).

Product Description: An off white to slight yellow powder.

Pharmacodynamics: Neomycin is a broad-spectrum bactericidal antibiotic with particular activity against certain members of the Enterobacteriaceae family. Its mode of action is at the ribosomal level. When administered orally, only a fraction (<5%) is absorbed systemically, the remainder remains as the active compound in the gastro-intestinal tract of the animal. Neomycin is not inactivated by enzymes or food. These pharmacological properties lead to neomycin being an efficacious antibiotic in the treatment of enteric infections caused by organisms susceptible to neomycin.

Pharmacokinetics: Approximately 3% of dose of neomycin is absorbed after oral or rectal (retention enema) administration, but this can be increased if gut motility is slowed or if the bowel wall is damaged. Therapeutic levels are not attained in the systemic circulation after oral administration. Orally administered neomycin is nearly all excreted unchanged in the feces.

Indications:

Cattle (excluding veal calves): For the treatment and control of colibacillosis (bacterial enteritis) caused by Escherichia coli susceptible to neomycin sulfate in cattle (excluding veal calves).

Sheep: For the treatment and control of colibacillosis (bacterial enteritis) caused by Escherichia coli susceptible to neomycin sulfate in sheep.

Swine: For the treatment and control of colibacillosis (bacterial enteritis) caused by Escherichia coli susceptible to neomycin sulfate in swine.

Goats: For the treatment and control of colibacillosis (bacterial enteritis) caused by Escherichia coli susceptible to neomycin sulfate in goats.

Turkeys (growing): For the control of mortality associated with E. coli organisms susceptible to neomycin sulfate in growing turkeys.

Recommended Dose:

Cattle (excluding veal calves): 10 mg of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram) in divided doses for a maximum of 14 days. Treatment should continue 24-48 hours beyond remission of disease symptoms, but not to exceed a total of 14 days.

Swine: 10 mg of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram) in divided doses for a maximum of 14 days. Treatment should continue 24-48 hours beyond remission of disease symptoms, but not to exceed a total of 14 days.

Sheep: 10 mg of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram) in divided doses for a maximum of 14 days. Treatment should continue 24-48 hours beyond remission of disease symptoms, but not to exceed a total of 14 days.

Goats: 10 mg of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram) in divided doses for a maximum of 14 days. Treatment should continue 24-48 hours beyond remission of disease symptoms, but not to exceed a total of 14 days.

Turkeys (growing): 10 mg of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram) for 5 days. Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 5 consecutive days.

Mode of Administration: By oral.

Contraindications: Do not use if there is known hypersensitivity to active ingredient.

Warnings and Precautions: Add to drinking water or milk. Not for use in liquid supplements. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian.

Interaction with Other Medicaments: None known.

Pregnancy and Lactation: None known.

Side Effects: Ototoxicity, nephrotoxicity.

Overdose and Treatment: None known.

Storage Conditions: Do not store above 25°C (controlled temperature room). Seal and store in a cool dry place.

Withdrawal Periods:

Cattle excluding veal calves: 1 day before slaughter

Sheep: 2 days before slaughter

Swine and goats: 3 days before slaughter

Turkey: zero days before slaughter

Packaging available: 500 g per bottle.

Manufactured by:

zoetis

Zoetis Suzhou Manufacturing Co. Ltd.

No 690, Jian Lin Road,
Suzhou New District,
Jiangsu 215151, P.R. China

Product Registration Holder:

Zoetis Malaysia Sdn. Bhd. (8898-V)
Lot 3.1, Level 3, Tower 7, Avenue 3, Bangsar South,
No. 8, Jalan Kerinchi, 59200, Kuala Lumpur, Malaysia.

Date of Revision: Dec 2018