

FEN-GUARD 8% POWDER

FOR ANIMAL USE ONLY

Product Name :

FEN-GUARD 8% POWDER

Name and Strength of Active Substance(s)

Each kg contains 80 g Fenbendazole

Product Description

FEN-GUARD 8% POWDER is a white to almost white powder.

Pharmacodynamics

Fenbendazole is an anthelmintic belonging to the benzimidazole-carbamate group. It acts by interfering with the energy metabolism of the nematode. Fenbendazole inhibits the polymerisation of tubulin to microtubules. This interferes with essential structural and functional properties of the cells of helminths, such as formation of the cytoskeleton, formation of the mitotic spindle and the uptake and intracellular transport of nutrients and metabolic products. Fenbendazole is effective against adult and immature stages. Fenbendazole has an ovicidal effect on nematode eggs.

Pharmacokinetics

After oral administration fenbendazole is only partially absorbed. Following absorption, fenbendazole is rapidly metabolised in the liver mainly to its sulphoxide (oxfendazole) and further to its sulphone (oxfendazole sulphone). Oxfendazole is the main component detected in plasma, accounting for about 2/3 of the total AUC (ie the sum of the AUC for fenbendazole, oxfendazole and oxfendazole sulphone). Fenbendazole and its metabolites are distributed throughout the body, reaching highest concentrations in the liver. The elimination of fenbendazole and its metabolites occurs primarily via the faeces and to a small extent in the urine.

Indication

Treatment of infections with the following parasites:

In pigs:

- Gastrointestinal roundworms: *Ascaris suum*, *Strongyloides ransomi*, *Oesophagostomum* spp.
- Lungworms: *Metastrongylus* spp.

In cattle:

- Gastrointestinal roundworms (adults and L4 larvae): *Ostertagia ostertagi* (including inhibited larvae) *Trichostrongylus* spp, *Cooperia onchophora*, *Nematodirus helvetianus*, *Bunostomum phlebotomum*, *Strongyloides papillosus*, *Oesophagostomum radiatum*.
- Lungworms (adults and L4 larvae): *Dictyocaulus viviparus*.

Recommended Dosage

Cattle : 7.5 mg fenbendazole per kg bodyweight as a single oral administration , 9.375 g of product per 100 kg body weight mixed with the food.

Pigs : 5 mg fenbendazole per kg bodyweight as a single oral administration , 6.25 g of product per 100 kg body weight mixed with the food.

Route of Administration

Orally via feed mixing

Contraindication

None.

Warnings and Precautions

- Special precautions for use in animals Body weight should be measured as accurately as possible before calculating the dose. Resistors all benzimidazoles can develop when frequent and repeated use of a control of this family.
- Special precautions to be taken by the person administering the veterinary medicinal product to animals Avoid direct contact with skin. Wash hands after use. Impervious gloves are recommended when using the drug. In case of accidental ingestion, seek medical advice immediately .

Interactions with Other Medicaments

None known.

Statement on usage during pregnancy and lactation

Can be used during pregnancy and lactation.

Adverse reactions

None known.

Overdose and Treatment

None known.

Withdrawal Period(s)

Meat and offal : 8 days. Milk: zero days.

Shelf-life

3 years from date of manufacture. Opened bag and reconstituted feed is recommended to be used within 24 hours.

Maximum Residual Limit (MRL)

Milk (cattle), muscle, kidney, fat (cattle and pig) : 100 µg/kg, Liver (cattle and pig) 500 µg/kg (Ref. REGOV Version 2-2009)

Storage Conditions

Store in a dry place below 30°C. Keep out of reach of children. Jauhkan ubat dari kanak-kanak.

Disposal of Containers

Emptied bag must be safely disposed in trash according to local government regulations.

Dosage Forms and packaging available

Available in 500g and 1 kg aluminium sachet and 25 kg PP paper bag.

Name and Address of manufacturer

Life Biopharma Sdn. Bhd.(546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2014-07-22