

Vetoquinol	Nom du produit : Fercobsang MY
Créé par : A.STIFANI	Référence article : 454251 0118 C L
le :	Dimensions : 160 x 120 mm
Visa :	1 couleur
	
281 U	

WITHDRAWAL PERIOD

Cattle, sheep, goats:

Meat and offals: 0 day.

Milk: 0 day.

Pigs:

Meat and offals: 0 day.

STORAGE CONDITIONS

Do not store above 25°C.

Shelf-life after first opening: 28 days.

Keep out of the reach and sight of children.

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIAL IF ANY

Ask your veterinary surgeon how to dispose of medicines no longer require. These measures should help to protect the environment.

DOSAGE FORMS AND PACKAGING

Amber glass vials of 100 mL, closed with sterilised chlorobutyl rubbers and oversealed with sterilised aluminium caps.

VETERINARY USE ONLY

REG. NUMBER N° **MAL14090137HA**

NAME & ADDRESS OF MANUFACTURER
VETOQUINOL S.A
 Magny-Vernois
 70204 Lure (France)

NAME & ADDRESS OF REGISTRATION HOLDER:

Gladron Chemicals Sdn. Bhd.
No.7, Jalan TP7, UEP Industrial Park, 40400 Shah Alam,
Selangor Darul Ehsan, Malaysia

Date of revision: March 2018.

Change of name and address of registration holder from Age D'or to Gladron Chemicals.



Magny-Vernois 70200 LURE (France)



FERCOBSANG® SOLUTION

INJECTABLE AND DRINKABLE SOLUTION

Reconstituent

COMPOSITION

Iron (as ammonium citrate) 1.7500 mg
 Cyanocobalamine (vitamin B12) 0.0250 mg
 Nicotinamide (vitamin PP) 20.0000 mg
 Cobalt (as digluconate) 0.0065 mg
 Benzyl alcohol (E1519) 20.8000 mg
 Excipient Q.S. 1 ml

100 ml vial of orange to brown solution

Intramuscular (I.M.), subcutaneous (S.C.) and oral route

PHARMACODYNAMICS

Iron, provided by the ferric ammonium citrate, plays an essential role in oxygen metabolism and in energy metabolism.

Cyanocobalamin (or vitamin B12) is a water soluble vitamin which plays a role in folate metabolism.

Nicotinamide (or vitamin PP) is also a water soluble vitamin. This is an active moiety of two co-enzymes which, in association with the apoenzymes, are involved in different stages of the degradation or synthesis of carbohydrates, fatty acids and amino acids. This molecule also plays a key role during terminal oxidative degradation of substrates in the citric acid cycle by fixing hydrogen during the terminal phase and by transporting it to the flavinic enzymes of the respiratory chain.

Cobalt is a trace element required for the activity of microflora in the rumen. In ruminants, cobalt is converted into vitamin B12 which, after absorption, participates in metabolism of propionic acid, the main energy source in these animals.

PHARMACOKINETICS

Iron, provided by ferric ammonium citrate, is absorbed into the duodenum and stored in the intestinal mucosa in the form of ferritin. The liver, spleen, bone marrow and muscles also store some of it.

Cyanocobalamin (or vitamin B12) is absorbed by the intestinal mucosa. During intramuscular or subcutaneous administration, the plasma peak is obtained in less than one hour. Cyanocobalamin is excreted in the bile.

Absorption of nicotinamide (or vitamin PP) occurs throughout the intestinal wall. It is rapid and complete. Distribution occurs rapidly through all tissues. Nicotinamide is transformed into an N-methyl derivative. Urinary excretion occurs in very small quantities in free form and mainly in metabolite form.

Cobalt gluconate is poorly absorbed in the gastrointestinal tract. The highest concentrations are observed in the liver, adrenals and thyroid. During parenteral administration, 90% of the cobalt is excreted in the urine within 48 hours. After ingestion, excretion occurs via the faecal route.

INDICATIONS

In cattle, sheep, goats, pigs, dogs and cats.

Treatment and prevention of deficiencies of iron, cobalt, vitamin B12 and vitamin PP (nicotinamide/vitamin B3), especially during:
 - convalescence from debilitating infectious or parasitic diseases
 - growth disorders
 - anemia.

RECOMMENDED DOSAGE / METHOD OF ADMINISTRATION

0.08 to 0.4 mg of iron, 1 to 5 µg of vitamin B12, 0.8 to 4 mg of vitamin PP and 0.27 to 1.4 µg of cobalt per kg of live weight per day depending on the species and in accordance with the following recommendations:

CATTLE: Intramuscular, subcutaneous and oral route:

- adult 20 ml
 - young 10 ml

SHEEP, GOATS: Intramuscular, subcutaneous and oral route:

- adult 5 ml
 - young from 1 to 2 ml

PIGS: Intramuscular, subcutaneous and oral route:

- adult 10 ml
 - fattening 5 ml

DOGS, CATS: oral route only:

1 to 2 drops per kg b.w.

The administration of the product should be made during 3-5 days.

CONTRA-INDICATIONS

Do not use in case of hypersensitivity to the active substances or to any of the excipients.

WARNINGS AND PRECAUTIONS

Special warnings for each target species : None.

Special precautions for use in animals : None.

Special precautions to be taken by the person administering the veterinary medicinal product to animals : None.

INTERACTIONS WITH OTHER MEDICAMENTS

None.

PREGNANCY AND LACTATION

Use of this drug will depend on the benefit/risk evaluation carried out by the veterinary surgeon.

ADVERSE EFFECTS / UNDESIRABLE EFFECTS

Not known.

OVERDOSE AND TREATMENT

Not known.

INCOMPATIBILITIES

Not Known.