

FOTAGEL Suspension 3g/20mL

Dioctahedral Smectite

NON-PRESCRIPTION

[Composition]

Each 20mL pouch contains 3g of dioctahedral smectite.

Other ingredients : D-sorbitol solution, Concentrated glycerin, Melon flavour S-J-AD (1501025), Vanilla flavour #V-21011, Potassium dihydrogen phosphate, Xanthan gum, Purified water

Preservatives :

- Methylparaben 0.001 % w/v
- Propylparaben 0.009 % w/v

[Description]

Vanilla and melon-flavoured light brown suspension contained in a Al-foil pouch.

[Pharmacodynamics]

Diosmectite (dioctahedral smectite) is a natural aluminomagnesium silicate taking the form of layers of fine sheet. It possesses a powerful coating property on the gastrointestinal mucosa due to its leaflet structure and its high plastic viscosity. Diosmectite increases the resistance of the mucosal gel in response to aggressive agents, by interacting with glycoproteins of mucus. By its action on the gastrointestinal mucous barrier and its high-binding capacity, diosmectite protects the gastrointestinal mucosa. It is radiolucent and does not colour the stools. At usual doses, it does not modify the physiological intestinal transit time.

[Pharmacokinetics]

Diosmectite is neither absorbed nor metabolized and it is eliminated through the faeces.

[Indications]

- Acute and chronic diarrhoea.
- Symptomatic treatment of pain associated with oesogastroduodenal and colic disease.

[Recommended Dosage]

Treatment of Acute Diarrhoea

Children :

- 2 years and older : 4 pouches per day for 3 days, then 2 pouches per day (up to maximum of 7 days).

Adults :

- An average of 3 pouches per day.
- In practice, the daily dosage can be doubled at the beginning of the treatment.

Other Indications

Children :

- 2 years and older : 2 - 3 pouches per day (up to maximum of 7 days).

Adults :

- An average of 3 pouches per day.

[Method of Administration]

Children :

Can be taken directly from the pouch, or added into feeding bottle and further dilute with water, to be given during the day, or mixed with a semi-liquid food: broth, stewed fruit, mashed vegetables or baby food.

Adults :

Can be taken directly from the pouch.

[Contraindications]

Hypersensitivity to diosmectite or to any of the excipients.

[Warnings and Precautions]

Diosmectite must be used with care in patients with a history of severe chronic constipation. In infants and children below 2 years, the use of FOTAGEL should be avoided. The reference treatment in acute diarrhoea is oral rehydration solution (ORS).

In children above 2 years, acute diarrhoea must be treated in conjunction with early administration of an oral rehydration solution (ORS), to avoid dehydration. The chronic use of FOTAGEL should be avoided.

In adults, the treatment does not dispense with rehydration, if this is considered to be necessary. The amount of rehydration by oral rehydration solution or intravenously must be adapted to the intensity of the diarrhoea, and the patient's age and characteristics.

If the symptoms do not improve within 7 days, consult your physician. If pain is associated with fever or vomiting, consult your physician.

[Interactions with Other Medicaments]

As the absorbent properties of this product may interfere with the rates and/or levels of absorption of other substances, it is recommended not to administer any other drugs at the same time.

[Pregnancy and Lactation]

Not recommended to be used during pregnancy or breastfeeding.

[Side Effects]

Very rare cases of development or aggravation of constipation have been reported; treatment was able to be continued after decreasing the dosage.

[Symptoms and Treatment of Overdose]

Symptoms of overdose include constipation, vomiting and bezoar.

In case of overdose, treatment should be decreased or discontinued, particularly in case of vomiting. Management should be symptomatic and supportive. If symptoms persist or worsen, the patient should be advised to consult a doctor.

[Storage Condition]

Store below 30°C.

Keep out of reach of children.

Jauhi daripada kanak-kanak.

Shelf life : 2 years

[Packing]

20mL suspension in one pouch, 20 pouches/box

[Revision Date] 22 April 2025

[Manufactured by]

Daewon Daewon Pharm.Co.,Ltd.

Head Office : 386, Cheonhodaero, Seongdong-gu, Seoul, Republic of Korea

Plant : 44, Yongso 2-gil, Gwanghyewon-myeon, Jincheon-gun, Chungcheongbuk-do, Republic of Korea