



SALOSONE OINTMENT

Description: White yellowish ointment

Content:

Betamethasone dipropionate (eq. to betamethasone 0.05%)

0.065% w/w

Salicylic acid

3.00% w/w

Indication:

It is used for the relief of seborrheic dermatitis, ichthyosis vulgaris, psoriasis and other inflammatory, corticosteroid-responsive dermatoses, hyperkeratotic and dry, scaling skin conditions.

Pharmacology:

Betamethasone dipropionate is a glucocorticoid applied topically in the management of various disorders in which corticosteroids are indicated. It has marked antipruritic, vasoconstrictor and anti-inflammatory activity.

Salicylic acid is a keratolytic with bacteriostatic and fungicidal actions.

Betamethasone dipropionate is well absorbed from the sites of local application, particularly under an occlusive dressing or when the skin is broken. It may be absorbed to give systemic effects. It is more slowly metabolized in the skin and this tends to be systemically absorbed to a greater extent. Salicylic acid is also rapidly absorbed through the skin.

Contraindications:

It is contraindicated in patients hypersensitive to salicylic acid and betamethasone dipropionate or any related components.

Precautions:

Steroids applied topically over large areas, particularly when the skin is broken, with or without occlusive dressings may result in systemic effects. Therefore, suitable precautions should be taken. Special precautions have to be observed when used on infants and children as it may cause growth retardation on prolonged use. Therefore, avoid prolonged use in children and on the face.

Its use should be discontinued if excessive dryness, irritation or sensitization of the skin occurs.

Avoid contact with the eyes and mucous membranes.

Use in pregnancy and lactation:

Risk benefits should be considered when using on pregnant or lactating mothers although safety of use has not been established.

Drug Interactions:

None known

Adverse effects:

When applied topically, particularly to large areas or when the skin is broken or under occlusive dressings, betamethasone dipropionate may be absorbed in sufficient amounts to cause systemic effects, eg. burning, itching irritation, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, hypertrichosis and folliculitis.

Dosage and administration:

Apply over affected skin twice daily

Treatment of over dosage:

Symptoms: Excessive or prolonged use of topical corticosteroids can suppress pituitary-adrenal function, resulting in secondary adrenal insufficiency, and produce manifestations of hypercorticism, including Cushing's disease. Prolonged use of salicylic acid causes salicylism.

Treatment: Symptomatic treatment is applied. Slow withdrawal of steroids is advised and treat electrolyte imbalance, if any. For salicylism, sodium bicarbonate is given orally to alkalinize urine and force diuresis.

Storage conditions : Keep container tightly closed. Store in a cool, dry place below 30°C.
Protect from light. Keep away from children.

Presentation : 15gm/ tube.

Product registration holder : PRIME PHARMACEUTICAL SDN. BHD.

& Manufacturer 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia

21/08/2025