

PABRON™ COUGH-D SYRUP

パブロン



Relieves Dry Cough & Nasal Congestion

Anti-Tussive & Nasal Decongestant
Alcohol-Free & Sugar-Free



**PABRON™
COUGH-D
SYRUP**



**PABRON™
COUGH-D
SYRUP**



**PABRON™
COUGH-D
SYRUP**



Each 5 ml syrup contains:

Tripolidine HCl	1.25mg
Pseudoephedrine HCl	30mg
Dextromethorphan HBr	15mg
Sodium Benzoate as Preservative	0.1% w/v

Product Description

A dark brown syrup with herbal flavour.

Pharmacodynamics

Tripolidine HCl has the actions of the histamine H1-receptor antagonists by diminishing or abolishing the main actions of the histamine in the body by competitive reversible blockade of histamine receptor sites on tissues. It is used for the symptomatic relief of common allergic symptoms of the respiratory tract. It has mild central nervous system depressant properties which may cause drowsiness.

It is used in combination with Pseudoephedrine HCl for rhinitis and in combination with other compound preparations for the symptomatic treatment of coughs and the common cold.

Pseudoephedrine HCl acts on alpha-adrenergic receptors in the mucosa layer of the respiratory tract, producing vasoconstriction. This shrinks swollen nasal mucous membrane, reduces tissue hyperemia, oedema and nasal congestion and increases nasal airway patency.

Dextromethorphan HBr is a cough suppressant which has a central action on the cough centre in the medulla. Dextromethorphan has very little sedative activity. It is used for the relief of unproductive cough and is reported to act within half an hour of administration by mouth and to exert for up to 6 hours.

Pharmacokinetics

After absorption from the gastrointestinal tract, Tripolidine HCl is metabolized into carboxylated derivative.

Pseudoephedrine is readily absorbed from the gastrointestinal tract. It is resistant to metabolism by monoamine oxidase and is largely excreted unchanged in the urine together with small amounts of its hepatic metabolite. It has a shelf-life of several hours.

Dextromethorphan is rapidly absorbed from the gastrointestinal tract. It is metabolized in the liver and excreted in the urine as unchanged dextromethorphan and demethylated metabolites including dextrothorphan, which has some cough suppressant activity.

Indications

Relieves dry cough & nasal congestion

Recommended Dosage

Adults & Children above 12 years:
10ml to be taken 3 times daily
Children between 6 to 12 years:
5ml to be taken 3 times daily
Children between 2 to 6 years:
2.5ml to be taken 3 times daily

Side Effects

Central nervous system depression or excitation may occur, drowsiness being reported most frequently. Sleep disturbance and rarely, hallucinations have been reported.

Skin rashes, with or without irritation, tachycardia, dryness of mouth, nose and throat have occasionally been reported. Urinary retention has been reported occasionally in male patients in whom prostatic enlargement could be an important predisposing factor.

Side effects attributed to dextromethorphan are uncommon; occasionally nausea, vomiting or gastrointestinal disturbance may occur.

Contraindications

Pabron Cough-D is contraindicated in individuals who have previously exhibited intolerance to it or to any of its constituents.

Pabron Cough-D is contraindicated in patients with severe hypertension or severe coronary artery disease.

Pabron Cough-D is contraindicated in individuals who are taking, or have taken, monoamine oxidase inhibitors within the preceding two weeks. The concomitant use of pseudoephedrine and this type of product may occasionally cause a rise in blood pressure.

The antibacterial agent, furazolidone, is known to cause a dose-related inhibition of monoamine oxidase. Although there are no reports of hypertensive crisis caused by the concurrent administration of Pabron Cough-D and furazolidone they should not be taken together.

Pabron Cough-D should not be administered to patients where cough is associated with asthma or where cough is accompanied by excessive secretions.

Dextromethorphan, in common with other centrally acting antitussive agents, should not be given to patients in, or at risk of developing respiratory failure.

Warning & Precautions

This medication may cause drowsiness and impair performance in test of auditory vigilance. If affected, do not drive or operate machinery.

Avoid alcohol or other centrally acting sedatives.

Use with caution for patients with hypertension, heart disease, diabetes, hyperthyroidism, elevated intra-ocular pressure, prostatic enlargement, renal and hepatic impairment.

Not to be used in children less than 2 years of age.

To be used with caution and on doctor's or pharmacist's advice in children 2 to 6 years of age.

Interactions with Other Medication

Concomitant use of Pabron Cough-D with sympathomimetic agents like decongestant, tricyclic antidepressant,

amphetamine-like psychostimulants and monoamine oxidase inhibitors may occasionally cause a rise in blood pressure.

The content of Pseudoephedrine may interfere with alpha- and beta-blocking drugs.

Symptoms & Treatment for Overdose

Overdose of Pabron Cough-D may cause drowsiness, dizziness, lethargy, ataxia, hyperactivity, respiratory depression, dryness of the skin and mucous membrane, tachycardia, hypertension, hyperpyrexia, irritability, convulsions, difficulty with micturition, nystagmus, nausea and vomiting.

Measures to maintain and support respiration and control convulsions, gastric lavage, catheterization of the bladder are necessary.

Should there be a need to eliminate Pseudoephedrine faster, dialysis or acid diuresis can be considered.

Naloxone can be considered to reverse the effect of Dextromethorphan.

Pregnancy & Lactation

Safe use in pregnancy has not been established.

Pseudoephedrine and Tripolidine are excreted in breast milk in small amounts but the effect of this on breast-fed infants is not known.

It is not known whether dextromethorphan or its metabolites are excreted in breast milk.

Route of Administration: Oral

Packing: 120ml in an amber PET bottle.

Storage

Store below 30°C.

Shelf-Life

2 years from the date of manufacture.

Manufactured by:

Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5, Cheng Industrial Estate,
75250 Melaka, Malaysia

Under Licence of:

Taisho Pharmaceutical Co., Ltd.
Tokyo, Japan

Product Registration Holder:

HOE Pharmaceuticals Sdn. Bhd.
Lot 10, Jalan Sultan Mohamed 6,
Bandar Sultan Suleiman,
42000 Port Klang,
Malaysia.

Revision date: May 2018

PA0800-002

