

Muposten Ointment 2%w/w

Description: A cream to light yellowish smooth semisolid mass

Composition: Each gram contains Mupirocin 20mg

Pharmacodynamics

Muposten ointment is a topical antibacterial agent, active against those organisms responsible for the majority of skin infections, eg *Staphylococcus aureus*, including methicillin-resistant strains, other staphylococci, and streptococci. It is also active against Gram-negative organisms such as *Escherichia coli* and *Haemophilus influenzae*.

Mupirocin potently inhibits bacterial protein and RNA synthesis by inhibition of isoleucyl-transfer RNA synthetase.

Pharmacokinetics

After topical application of Muposten ointment, mupirocin is only very minimally absorbed systemically and that which is absorbed is rapidly metabolised to the antimicrobially inactive metabolite, monic acid. Penetration of mupirocin into the deeper epidermal and dermal layers of the skin is enhanced in traumatized skin and under occlusive dressing.

Indications

Treatment of bacterial skin infections, e.g. impetigo, folliculitis and furunculosis.

Recommended dose

Muposten ointment should be applied to the affected area up to 3 times daily, for up to 10 days. The area may be covered with a dressing or occluded if desired.

Contraindications

Hypersensitivity to Muposten ointment or other ointments containing polyethylene glycol.

Warnings and Precautions

When Muposten ointment is used on the face, care should be taken to avoid the eyes.

Polyethylene glycol can be absorbed from open wounds and damaged skin and is excreted by the kidneys. In common with other polyethylene glycol-based ointment, Muposten ointment should be used with caution if there is evidence of moderate or severe renal impairment.

In the rare event of a possible sensitization reaction or severe local irritation occurring with the use of Muposten ointment, treatment should be discontinued, the product should be rinsed off and appropriate alternative therapy for the infection instituted.

Interaction with other Medicaments

None stated.

Pregnancy and Lactation

Studies in experimental animals have shown mupirocin to be without teratogenic effects. However, there is inadequate evidence of safety to recommend the use during pregnancy or lactation.

Side Effects

Muposten ointment is well tolerated. The following side effects have been reported very rarely: itching, burning, erythema, stinging and dryness localized to the area of application. Cutaneous sensitization reactions to mupirocin or the ointment base have been reported rarely. Systemic allergic reactions have been reported with Muposten ointment.

Symptoms and Treatment overdose

The toxicity of mupirocin is very low. In the event of overdose, symptomatic treatment should be given.

Storage Condition

Store below 30°C. Shelf life: 3 years

Presentation

Aluminum tubes of 3.5g, 5.0g, 10.0g and 15.0g.

Manufactured by:

Noripharma Sdn. Bhd., Lot 5030, Jalan Teratai, 5 ½ Mile off Jalan Meru, 41050 Klang, Selangor.

Date of Revision: December 2013