

SAME SIZE ARTWORK
LEAFLET SIZE: 88 mm x 186 mm

24 mm

SPACE FOR PHARMACODE

For the use only of a Registered Medical Practitioner
or a Hospital or a Laboratory

SaferonTM
Chewable Tablets

SPACE FOR PHARMACODE

Composition
Each uncoated chewable tablet contains
Iron (III)-Hydroxide polymaltose complex
equivalent to Elemental Iron 100 mg
Folic Acid BP 500 mcg

Preservatives :
Methyl Hydroxybenzoate BP 0.9 mg
Propyl Hydroxybenzoate BP 0.09 mg

Description
Brown coloured speckled, round biconvex, uncoated tablets having
chocolate flavour and sweet taste.

Indications
Treatment of latent iron deficiency and iron deficiency anaemia.

Pharmacodynamics
Iron (III) hydroxide polymaltose complex is non-toxic, due to the fact
that an active transport of iron or a rate determining competitive
ligand exchange instead of passive diffusion process is assumed to
occur. In contrast, the absorption of Ferrous salts is not controlled by
or a respective feed back mechanism, even in normal or iron-loaded
subjects, and therefore, adverse drug reactions may occur
especially when administering several doses a day.

Dosage & Administration
Dosage and duration of therapy are dependent upon the extent of
iron deficiency. The daily dose can be divided into separate doses or
can be taken at one time. Saferon chewable tablets can be chewed
or swallowed whole and should be taken during or immediately after
a meal. In case of manifest iron deficiency the therapy takes 3-5
months until a normalisation of the haemoglobin value is achieved.
Afterwards the therapy should be continued for several weeks with a
dosage such as described for latent iron deficiency to replenish the
iron stores.

	Manifestation of iron deficiency	Latent iron deficiency
Saferon	1-3 tablets daily	1 tablet daily

Contraindications
Hypersensitivity to any ingredient of the formulation.

Precautions
Saferon contains aspartame and may not be suitable for
phenylketourics.

Pregnancy and Lactation
Saferon Chewable Tablets should not be administered in the first
trimester of pregnancy. Saferon Chewable Tablets should only be

PE00000 MY

186 mm

88 mm

SAME SIZE ARTWORK
LEAFLET SIZE: 88 mm x 186 mm

186 mm

administered in the second and third trimester of pregnancy if the
benefits of treatment outweigh the potential risk to the foetus. No
controlled studies are available on animals or on pregnant women.

It is not known whether the Iron Polymaltose Complex is excreted in
human milk. A decision should be made whether to discontinue
nursing or to discontinue the drug, taking into account the
importance of the drug to the mother.

Side Effects
Rarely nausea, vomiting, diarrhea & constipation occurred.

Interactions
No interactions have been observed. Since, the iron is complex
bound, ionic interactions with foodstuff components (phytin,
oxalates, tannin, etc.) and concomitant administration of
medicaments (tetracyclines, antacids) are unlikely to occur. Saferon
does not cause teeth staining. The haemoccult-test (selective for
haemoglobin) for the detection of occult blood is not impaired and
therefore iron therapy must not be interrupted.

Effects on ability to drive and use machines
None.

Overdosage
In case of overdose neither intoxication nor iron overload have been
reported.

Incompatibilities
None known.

Storage
Store below 30°C. Protect from moisture & light.
Keep medicine out of reach of children.

Presentation
A box of 3 Strips of 10 Tablets each.
MAL06030848 X

Date of Revision: 11 October 2021

Product Registration Holder:
Glenmark Pharmaceuticals (Malaysia)
Sdn Bhd (660397-W)
D-31-02, Menara Suezcap 1,
No. 2, Jalan Kerinchi,
59200 Kuala Lumpur, Malaysia

Manufactured by:
 **glenmark**
PHARMACEUTICALS LTD.
Plot No. E-37, 39, D-Road, MIDC,
Satpur, Nasik - 422 007,
Maharashtra State, India.
© Trade Mark

PE00000 MY

88 mm

ICONGRAPHICS CODE:

PANTONE SHADE


PANTONE
BLACK
PROCESS C

**Supersedes
Artwork Code
PE9034**

PHARMACODE: