



BEVARATE-N CREAM

Description: White colour cream

Content: Betamethasone 17-valerate (eq. to betamethasone 0.1%)	0.12%w/w
Neomycin sulphate	0.50%w/w

Indication: Dermatitis (including seborrheic dermatitis, lichen simplex and lichen planus), eczema (including atopic and discoid eczema), psoriasis, insect bites, anal/ genital intertrigo and other inflammatory skin conditions, where corticosteroid is indicated and bacterial infection is present or suspected.

Pharmacology: Betamethasone valerate, a potent adrenocorticoid, diffuses across membranes and stimulates transcriptions of mRNA resulting in protein synthesis of various enzymes responsible for anti-inflammatory effects. Neomycin, an aminoglycoside antibiotic, bactericidal against common skin infection bacterias.

Betamethasone is absorbed percutaneously, Occlusion, use over extensive areas, or prolonged use increases topical absorption. It is bound to plasma proteins, metabolized in the liver and excreted in kidneys. Once neomycin is absorbed, it is rapidly and completely excreted in the urine in its active form.

Contraindications: Patients hypersensitive to neomycin, betamethasone or any related components, otitis externa, prolonged use on large areas, acne vulgaris, rosacea, perioral dermatitis, infants <2yrs, viral infections and perianal/genital pruritis.

Precautions: Steroids applied over large areas, particularly on broken skin, with/out occlusive dressings result in systemic effects. Precautions has to be observed when use on infants & children as it may cause growth retardation on prolonged use. Avoid prolonged use on the face. Flush with water if the cream enters the eye. The topical use of neomycin in patients with extensive skin damage or perforated tympanic membranes may result in deafness. Prolonged local use should be avoided as it may lead to skin sensitization.

Use in pregnancy and lactation:

It is not recommended because of neomycin causing theoretical risk of foetal toxicity, unless directed by doctor.

Drug Interactions:

Neuromuscular blocking agents - neomycin sulphate intensifies and prolongs respiratory depressant effects.

Side effects: Side effects may be pronounced with use of occlusive dressing. With prolonged use, moon face, thinning of skin with easy bruising, acne, facial hair growth, loss of hair, Cushing's syndrome and striae may occur. Hypersensitivity reaction, usually of delayed type, occur frequently during local treatment with neomycin but may be masked by the combined use of steroids. Cross sensitivity with aminoglycoside antibiotics may occur.

Dosage and administration : Apply sparingly to affected area 2 - 3 times dly, Upon improvement, application is reduced to once dly or less. Duration is less than 7 days unless under medical supervision. For children, duration is less than 5 days.

Treatment of over dosage: Acute overdosage is unlikely to occur, however, in case chronic overdosage or misuse, the features of hypercorticism may appear, in this situation, topical steroids should be discontinued.

Storage conditions : Keep container tightly closed. Store in a dry place at below 30°C. Protect from light. Keep away from children.

Shelf life : 3 years

Presentation : 15gm/ tube

Manufactured & : **PRIME PHARMACEUTICAL SDN. BHD.**

Distributed by : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah, 14000 Bukit Mertajam, Penang, Malaysia.