

# DOPROCT SUPPOSITORY & DOPROCT RECTAL OINTMENT

## Content:

Each suppository contains: hydrocortisone acetate 7.5mg, benzocaine 40mg and zinc oxide 250mg.

Each gram of ointment contains hydrocortisone acetate 5.0mg, benzocaine 10mg & zinc oxide 200mg

## Description:

**Doproct Suppository:** White, oblong-shape suppository.

**Doproct Rectal Ointment:** White ointment

## Properties

Hydrocortisone is produced in the adrenal cortex and belongs to the mild group of corticosteroids with anti-inflammatory and anti-pruritic actions.

Benzocaine is a local anaesthetic used for surface anaesthesia to block the sensory nerve endings in the skin or mucous membranes.

Zinc oxide is mildly astringent and is used topically as a soothing and protective application for haemorrhoids.

**Doproct** is a combination of hydrocortisone acetate, benzocaine and zinc oxide. The combined properties of these agents break the itch-scratch-cycle, thus allowing healing of the condition.

Less than 50% of hydrocortisone is absorbed following rectal application. Corticosteroids in the circulation are extensively bound to plasma protein, mainly globulin and less to albumin. Only unbound hydrocortisone has pharmacological effects or is metabolised. Corticosteroids are metabolised mainly in the liver but also in the kidney, and are excreted in the urine.

Benzocaine is minimally absorbed and therefore relatively free from systemic side effects or adverse drug interactions. Onset of action is rapid, with initial effects obtained in about 1 minute and action lasting about 15 - 20 minutes. It is metabolised hepatically and the metabolites are renally excreted.

## Indication

For treatment of pain, itching and inflammation arising from irritated anorectal tissues e.g. anal pruritus, proctitis, haemorrhoids, fissures and postoperative haemorrhoidectomy.

## Recommended Dosage

By rectal route only.

**Doproct Suppository:** Insert 1 suppository in the anal canal 2 times daily in the morning and evening after bowel movement. This product is best used with **Doproct Rectal Ointment**.

**Doproct Rectal Ointment:** Applied a small quantity to the anal canal 2 times daily in the morning and evening after each stool. The ointment may be used separately or concurrently with Doproct Suppositories.

## Contraindications

**Doproct** is contraindicated in patients having a history of hypersensitivity to hydrocortisone acetate, benzocaine or any of its ingredients.

## Warnings / Precautions

**Doproct** should be used with caution in patients with traumatised mucosa or sepsis in anorectal region.

**Doproct** should not be used until an adequate proctological examination is completed and malignancy excluded. Other specific measures against infections, allergy, and other causal factors must not be neglected.

When administering **Doproct** to the rectum, care should be taken to avoid the introduction of an excessive amount. It should be borne in mind that the systemic availability is relatively high with rectal administration, and large doses can cause effects on the central nervous system. Prolonged use of large doses of hydrocortisone can result in systemic effects or local effects such as skin atrophy. Patients should be advised to inform physicians of the previous use of hydrocortisone. In the event of infections caused by viruses, bacteria, pathogenic fungi or parasites, local glucocorticoids must not be used without concomitant causal therapy.

If irritation or rectal bleeding develops, the product should be discontinued, the patients examined and appropriate therapy instituted.

## Interaction with Other Medicaments

Ester derivatives such as benzocaine may antagonise the activity of aminosalicyclic acid or sulfonamides.

## Pregnancy and Lactation

Hydrocortisone acetate is excreted into breast milk in such small quantities that any effect on the nursing infants seems unlikely at therapeutic doses. However, as the safety of topical corticosteroids during pregnancy has not been fully established, it should not be used on extended areas, in large amounts or for prolonged periods of time.

## Side Effects

Contact sensitivity such as dry skin, pruritus, irritation, transient burning or stinging sensations may occur.

## Symptom & Treatment of Overdose

When used as indicated, contact sensitivity such as dry skin, pruritus, irritation, transient burning or stinging sensations may occur if **Doproct** is used excessively. Discontinue use if such symptoms occur.

When swallowed, an acute single overdose of **Doproct** does not produce toxic effects. Treatment of a large acute overdose should include gastric lavage, purgation with magnesium sulphate and complete bed rest. If necessary, give oxygen and general supportive measures.

## Storage Condition

Keep in a cool dry place, protect from light.

Keep out of the reach of children.

*Jauhi daripada kanak-kanak.*

**Doproct Suppository:** Store at 2 - 8 °C.

**Doproct Rectal Ointment:** Store below 30°C.

**Shelf life:** Please refer to outer box label

## PACKING / PACK SIZES

**Doproct Suppository:**

PVC/PE white film-strip containing 5 suppositories per strip.

Box of 2 x 5 suppositories

**Doproct Rectal Ointment:**

Aluminium tube of 10g.

Box of [10 x 10g] tubes.

## Manufactured by:

Continental-Pharma Co. Ltd.  
Bangkok, Thailand.

Revised Date: Nov 2019