

PACKAGE INSERT

PROZINE 10 SYRUP 10MG/5ML

Each 5ml contains:
Promethazine HCl 10mg

Product Description: A clear, brown syrup with blackcurrant flavour.

Pharmacodynamics:
Promethazine hydrochloride, a phenothiazine derivative, possesses anti-histamine, sedative, anti-motion sickness, anti-emetic, and anti-cholinergic effects. The duration of action is generally from four to six hours. The major side reaction of this drug is sedation. As an anti-histamine, it acts by competitive antagonism but does not block the release of histamine. It antagonises in varying degree, most but not all of the pharmacological effects of histamine.

Pharmacokinetics:
It is readily absorbed from the gastro-intestinal tract. Following oral administration, the effects start within 15 to 20 minutes and are fully developed within one hour. The highest concentration occurs in the lungs, progressively lower concentrations in the spleen, kidney, brain, muscle, and skin. Small amounts of the drug are excreted unchanged, most appear as degradation products that are almost completely excreted within 24 hours. The main site of metabolic transformation is the liver, but the lungs and kidneys can also metabolise the compound.

Indication:
In addition to its antihistaminic action, which is of marked potency and prolonged duration, it also provides antiemetic as well as sedative actions. Promethazine HCl is useful in the management of the following conditions:
a) Dermographism
b) Vasomotor rhinitis
c) Perennial and seasonal allergic rhinitis
d) Antiemetic therapy in postoperative patients
e) Preoperative, post-operative or obstetric sedation
f) Amelioration of allergic reactions to blood or plasma
g) Active and prophylactic treatment of motion sickness
h) Allergic conjunctivitis due to inhalant allergens and foods
i) Mild, uncomplicated allergic skin manifestation of urticaria and angioedema
j) Prevention and control of nausea and vomiting associated with certain types of anaesthesia and surgery
k) Sedation in both children and adults as well as relief of apprehension and production of light sleep from which the patient can be easily aroused

Recommended Dosage:
Adults:
The average oral dose is 25 mg, taken before retiring; however, 12.5 mg may be taken before meals and on retiring, if necessary. For children more than 2 years, a single 25 mg dose at bedtime or 5 mg to 15 mg taken three times daily will usually suffice.
After initiation of treatment in children (more than 2 years) or adults, dosage should be adjusted to the smallest amount adequate to relieve symptoms.
Motion sickness:
The average adult dose is 25 mg taken twice daily. The initial dose should be taken one-half to one hour before anticipated travel and be repeated eight to twelve hours later, if necessary. On succeeding days of travel, it is recommended that 25 mg be given on arising and again before the evening meal.
For children (more than 2 years), 12.5 mg to 25 mg twice daily may be administered.
For nausea and vomiting in children the dose should be adjusted to the age and weight of the patient and the severity of the condition being treated.
For prophylaxis of nausea and vomiting, as during surgery and the post-operative period, the average dose is 25mg repeated at four to six hour intervals as necessary.
Sedation:
Administration of 12.5 mg to 25 mg induces a quiet sleep from which the patient can be easily aroused. Adults usually require 25 mg for night-time, pre-surgical, or obstetrical sedation.

Route of Administration: Oral

Contraindications:
Promethazine is contraindicated in comatose states, in patients who have received large amounts of central nervous system depressants (e.g. alcohol, barbiturates, narcotics, etc.), and in patients who have demonstrated an idiosyncrasy or hypersensitivity to promethazine. It is contraindicated for use in the treatment of lower respiratory tract symptoms, including asthma.

Warnings and Precautions:
Promethazine should be used with caution in patients with narrow-angle glaucoma, prostatic hypertrophy, stenosing peptic ulcer, pyloroduodenal obstruction, and bladder-neck obstruction. It should be used with caution in patients with bone marrow depression. Ambulatory patients should be cautioned against driving automobiles or operating dangerous machinery until it is known that they do not become drowsy or dizzy from promethazine hydrochloride therapy.
Children: Should not be used for children under 2 years of age as safety is not established.
Excessively large dosages of promethazine in children may cause hallucinations, convulsions, and sudden death.
• Promethazine should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression.
• To be used with caution and under doctor's/ pharmacist's advice in children 2 to 6 years of age.
Anti-emetics may mask the symptoms of an unrecognised disease and thereby interfere with diagnosis.

Interactions with other Medicaments:
The sedative action of promethazine hydrochloride is additive to the sedative effects of CNS depressants; Therefore, agents such as alcohol, barbiturates, and narcotic analgesics should either be eliminated or given in reduced dosage in the presence of promethazine hydrochloride. When given concomitantly with promethazine hydrochloride, the dose of barbiturates should be reduced by at least one-half, and the dose of analgesic depressants, such as morphine or meperidine, should be reduced by one-quarter to one half.

Pregnancy and Lactation:
The safe use of promethazine has not been established with respect to the possible adverse effects upon fetal development. Therefore, the need for the use of this drug during pregnancy should be weighed against the possible but unknown hazards to the developing fetus.
It is not known whether promethazine is excreted in human milk. Caution should be exercised when promethazine is administered to a nursing mother.

Side Effects:
Autonomic reactions such as dryness of the mouth, blurring of vision, and rarely dizziness may occasionally arise. Very rare cases have been reported where patients receiving promethazine have developed leukopenia. In one instance, agranulocytosis has been reported. In nearly every instance reported, other toxic agents known to have cause these conditions have been associated with the administration of promethazine.
Cardiovascular effects from promethazine have been rare. Minor increases in blood pressure and occasional mild hypotension have been reported. Photosensitivity, although extremely rare, has also been reported. Occurrence of photosensitivity may be a contraindication to further treatment with promethazine or related drugs.
Suicidal doses have resulted in deep sedation, coma, and rarely convulsions and cardiorespiratory symptoms compatible with the depth of sedation present.
A paradoxical reaction has been reported in children receiving single doses of 75mg to 125mg orally, characterised by hyperexcitability and nightmares.

Symptoms and Treatment of Overdose:
Signs and symptoms of overdose range from mild depression of the CNS and cardiovascular system to profound hypotension, respiratory depression, and unconsciousness. Stimulation may be evident, especially in children and geriatric patients. Atropine-like signs and symptoms such as dry mouth, fixed, dilated pupils, flushing etc. as well as gastrointestinal symptoms may occur.
Treatment: Treatment of overdose is essentially symptomatic and supportive.
Early gastric lavage may be beneficial. Severe hypotension usually responds to the administration of levaterenol or phenylephrine. Extrapyramidal reactions may be treated with anticholinergics, anti-Parkinson agents, diphenhydramine, or barbiturates. Additional measures include oxygen and intravenous fluids.

Storage Conditions: Store below 30°C.

Pack Size: A bottle of 60ml, 100ml, 120ml.
Pack Size (export only): 1 litre.

Shelf-life: 3 years.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST .

Manufacturer & Product Registration Holder:
SUNWARD PHARMACEUTICAL SDN. BHD.
No. 9, 11 & 17, Jalan Kempas 4,
Taman Perindustrian Tampoi Indah,
81200 Johor Bahru, Johor, Malaysia.
SL327A-R1
Revision Date: 31/12/2025