



NORIT CAPSULE 200 MG
ACTIVATED CHARCOAL

COMPOSITION: Each capsule contains 200 mg Activated Charcoal as active ingredient.

PRODUCT DESCRIPTION: Norit capsules contains activated charcoal which is a fine, black, insoluble powder without taste or odor in black hard gelatin capsule that has a unique shorter and wider shape.

ROUTE OF ADMINISTRATION: Oral

PHARMACODYNAMICS:

Mechanism of Action: Activated charcoal is obtained by destructive distillation of organic material, coconut shells (which contain high carbon content and low ash and tar content), which are then 'activated' by exposing it to oxidizing STEAM at approximately 1000°C high temperature (in depleted oxygen environment), resulting in the production of vastly increased surface area from the creation of pores (with controlled pore size). The drug has the ability to adsorb a wide variety of substances onto its surface. This property can be applied in preventing the absorption of various harmful substances, which are responsible for diarrhea cases (traveler's disease), from the gastrointestinal tract and various drugs and toxicants involved with oral poisonings. Norit activated charcoal has fulfilled the requirements of **USA/ European Pharmacopoeia** for its adsorption capacity of **≥38% vs methylene blue** and **≥ 40% phenazone** respectively (the references test compounds in both respective pharmacopoeia).

Adsorption of substances onto charcoal is a reversible process with rapid adsorption and slow resorption. The rate of adsorption depends on the particle size, viscosity of the medium, pH (acids and bases) and the amount and type of gastrointestinal contents.

PHARMACOKINETICS:

Since activated charcoal is not absorbed from the gastrointestinal tract, there is no distribution phase nor any metabolism. The complex between activated carbon and the adsorbed compounds may only be passed in the stool. Norit activated charcoal is compliance with ICH Q3D guidelines on elemental impurities.

INDICATIONS:

Norit is indicated as gastrointestinal absorbent in the treatment of acute diarrhea, flatulence (excessive gas in the stomach and/or intestine) and mild food poisoning.

DOSAGE:

Diarrhea & Flatulence: 3 capsules 3-4 times a day; 12 capsules is the maximum dose per day. Children's dosage is half of adult's.

Mild food poisoning: 3 to 5 capsules repeated 3 to 4 times daily. Children's dosage is half dosage of adult dose.

The capsules should be taken with water orally.

Poison Information Centre (USM Penang)
Call-line: 1-800-88 8099/+6012-4309499

DRUG INTERACTIONS:

The use of activated charcoal will generally reduce the effect of oral medicines. Activated charcoal has the potential to reduce the absorption of many drugs from the gastrointestinal tract and simultaneous oral therapy should therefore be avoided.

To prevent this interaction, take activated charcoal at least 2 hours after medication taken orally.

CONTRAINDICATIONS:

Total gastro intestinal obstruction, suspected ileus, acute abdominal pain and ulcerative colitis.

WARNINGS & PRECAUTIONS:

- 1) The use of activated charcoal will cause the feces to be black.
- 2) Diarrhea: In case of persisting diarrhea after three days of treatment, consult a doctor for additional clinical and biochemical investigations.
- 3) Oral rehydration therapy which is the use of appropriate fluids including oral rehydration salts remains the most effective treatment for dehydration due to diarrhea. The intake of as much of these fluids as possible is therefore imperative.
- 4) Treatment of diarrhea in children under 6 years of age should be under medical supervision.

178.062/02-22R

PREGNANCY & LACTATION:

This medicine can, as far as is known, be used as prescribed during pregnancy and breastfeeding without any danger for the foetus or the child.

SIDE EFFECT

Activated charcoal is relatively non-toxic when given by mouth but gastrointestinal disturbances such as vomiting and constipation have been reported.

It may colour the faeces black.

SYMPTOMS & TREATMENT OF OVERDOSE:

Higher dosage or long time administration may cause nausea and constipation requiring the administration of laxative.

STORAGE CONDITION:

Store in the original package, in order to protect from moisture. Keep away from direct sunlight and store at temperature between 15-30 °C

SHELF LIFE:

5 years at 15-30 °C stored in PVC / blister strips after date of production.

DOSAGE FORMS AND PACKAGING AVAILABLE:

30 capsules per box

Manufactured by:

Norit Nederland B.V.
Mr. Ovingkanaal O.Z. 3,
7891 EV Klazienaveen,
The Netherlands

Product Registration Holder:

RX Pharma Sdn Bhd
No. 35, Pusat Pedada,
Jalan Pedada, 96000
Sibu Sarawak

Imported & Distributed by:

Ubisson Sdn. Bhd. (585048-H)
No. 35, 1st & 3rd Floor, Pusat Pedada,
Jalan Pedada, 96000 Sibu, Sarawak, Malaysia.
Tel: +6084-333858
Fax: +6084-311688
Email: office.ubisson@gmail.com

MAL REG. NO.: MAL06121041XZ

Revised Date: March 2022

Printproof

ATTENTION!

We need an approved proof before we can start the production.
Please fill in this form in clear writing.

Date:	02-08-2022	Proof no.: 4
Order no.:	415261	
Item no. Code:	178.062_02-22 Norit Capsules Maleisië	
Article no. Dobber:	N09718	
Size:	148 x 210 mm	DOBBER HEALTHCARE
Folded:	148 x 33 mm	LEAFLETS
Print color(s)		Dobber Healthcare Leaflets BV
Front:	PMS 286	Westerwerf 13
Back:	PMS 286	1911 JA Uitgeest
Square (oversteek):	-	Phone +31 (0)251 31 19 50
Optical code:	-	E-mail: info@dobberpharma.com
Pharma code:	-	www.dobberhealthcare.com
Font:	Helvetica Neue	
Font size:	8,5 pt	
Leading:	8,5 pt	

To be filled in by client

Approved: ☐ YES ☐ NO

Date:

Name and signature:

(This form belongs to SOP 'Werksaamheden DTP.Studio'.)



Proefsticker DHL (Mac/Server/Proefsticker DHL 14122018,12)