

Package Insert

Nervit B Forte Tablet

Composition:

Each tablet contains:

Thiamine Mononitrate (Vitamin B ₁)	250 mg
Pyridoxine HCl (Vitamin B ₆)	250 mg
Cyanocobalamin (Vitamin B ₁₂)	1 mg

Description:

A pink color round shaped film coated tablet with 13mm diameter.

Pharmacology:

Thiamine is a water-soluble vitamin. It is an essential nutrient required for carbohydrate metabolism and is also involved in nerve transmission. Pyridoxine, a water-soluble vitamin, is involved principally in amino acid metabolism, but is also involved in carbohydrate and fat metabolism. It is also required for the formation of haemoglobin. Cyanocobalamin influences hematopoiesis. It is involved in formation and metabolism of purines and pyrimidine, and in the synthesis of nucleoproteins.

Pharmacokinetics:

Thiamine, Pyridoxine and Cyanocobalamin are absorbed from gastrointestinal tract.

Thiamine is widely distributed to most body tissue; it is not stored in the body. Thiamine is excreted in urine as unchanged form or as its metabolites, pyrimidine.

Pyridoxine is converted to its active form pyridoxal phosphate and transformed to pyridoxic acid which is excreted in the urine.

Cyanocobalamin is bound to plasma proteins and stored in the liver. It is excreted in the bile and undergoes some enterohepatic recycling. It is also excreted in urine.

Indication:

For neurological and other disorders associated with disturbance of the nerve cell metabolism in which high-dose B complex vitamins play a role. These include: polyneuritis (of toxic and non-toxic etiology); neuralgias eg, lumbalgia, trigeminal neuralgia, shoulder-arm syndrome, herpes zoster, cervical syndrome, ischialgia, etc; diabetic neuropathy; metabolic and neuropathic changes due to pregnancy and oral contraceptives; convalescence.

Contraindication:

Known hypersensitivity to vitamins B₁, B₆ or B₁₂.

Should not be used in patients on Levodopa therapy.

Dosage and Directions:

Therapeutic Dose: Take 2-3 tablets daily after meals for 7-10 days.

Maintenance Dose: 1 tablet daily after meal.

Do not take the tablets on an empty stomach.

Warning & Precaution:

Cyanocobalamin should not be given before a diagnosis has been fully established because of the possibility of masking symptoms of subacute degeneration of the spinal cord.

Cyanocobalamin is not a suitable form of vitamin B₁₂ for the treatment of optic neuropathies associated with raised plasma concentrations of cyanocobalamin.

Long term administration of large doses of pyridoxine is associated with the development of severe peripheral neuropathies. This may occur with doses in excess of about 2g daily. Cyanocobalamin should not be used in megaloblastic anaemia of pregnancy. Administration of doses greater than 10 microgram daily may produce a haematological response in patients with folate deficiency.

Pregnancy and Lactation:

If you are pregnant or breastfeeding, please consult your doctor or healthcare professional before taking this product.

Thiamine is widely distributed to most body tissues and appears in breast milk. Pyridoxal and cyanocobalamin crosses the placenta and also appears in breast milk.

Adverse Reaction:

Allergic hypersensitivity reactions have occurred rarely following parenteral administration of vitamin B₁₂ compounds. Adverse effects seldom occur following administration of vitamin B₁, but hypersensitivity reactions have occurred, this is mainly after parenteral administration. Long-term administration of large doses of pyridoxine (in excess of 2g daily) is associated with the development of severe peripheral neuropathies

Interaction:

Loop diuretics (furosemide, ethacrynic acid, bumetanide): Chronic use of loop diuretics may result in thiamin deficiency. Concomitant intake of thiamin with food that contains sulfites may inactivate thiamin. Concomitant intake of thiamin with tea, coffee and decaffeinated coffee may inactivate thiamin. This can be reversed by taking supplements.

Pyridoxine reduces the effects of levodopa, but this does not occur if a dopa decarboxylase inhibitor is also given. Pyridoxine reduces the activity of almitriptan. It has also been reported to decrease serum concentrations of Phenobarbital and phenytoin. Many drugs may increase the requirement of pyridoxine; such drugs include hydralazine, isoniazid, penicillamine, and oral contraceptives.

Absorption of vitamin B₁₂ from the gastrointestinal tract may be reduced by neomycin, aminosalicylic acid, histamine H₂-antagonists, metformin, para-aminosalicylic acid, potassium chloride, colchicines, and proton pump inhibitors (lansoprazole, omeprazole, pantoprazole, rabeprazole). Cholestyramine and colestipol may decrease the enterohepatic reabsorption of vitamin B₁₂. Inhalation of the anesthetic agent nitrous oxide can produce a functional vitamin B₁₂ deficiency. The use of antibiotics may alter the intestinal microflora and may decrease the possible contribution of vitamin B₁₂ by certain inhabitants of microflora to the body's requirement for the vitamin.

Overdosage:

Treatment is symptomatic and supportive.

Storage Condition:

Store in a dry place below 30°C.

Shelf life:

2 years

Presentation:

Blister of 10 capsules. Box of 1, 3, 6, 10 and 50 blisters.

Manufacturer:

Pahang Pharmacy Sdn. Bhd. (10826-K)
Lot 5979, Jalan Teratai, 5 ½ Miles Off Jalan Meru,
41050 Klang, Selangor,
Malaysia.

Date revised: 091214