

FUCID CREAM 2% W/W

Composition:

Each gram of cream contains: Fusidic acid (Micronised) 20mg or 2.0% w/w

Preservative: Chlorocresol 1mg or 0.10% w/w

Description:

White to off white, smooth cream.

Pharmacodynamics:

Fusidic acid is an antibiotic with a bacteriostatic or bactericidal activity, mainly against Gram positive bacteria.

Staphylococci, including the strains resistant to penicillin or to other antibiotics, are particularly susceptible to fusidic acid. Besides pronounced antibacterial activity against the majority of organisms responsible for skin infections, it also possesses unique ability to penetrate to the focus of infection, even when applied to intact skin.

Pharmacokinetics:

Fusidic acid penetrates normal or inflamed skin and achieves bactericidal concentrations down to the lower corium.

Fusidic acid is excreted mainly in the bile with little excreted in the urine.

Indication:

Fusidic acid is indicated for skin infections caused by Staphylococci, Streptococci, *Propionibacterium acnes*, *Corynebacterium minutissimum* and other fusidic acid-sensitive organisms eg, impetigo, infected wounds, folliculitis, boils, sycosis barbae, carbuncles, hidradenitis, paronychia and erythrasma.

Recommended dosage:

Apply to the affected area 2-3 times a day, generally for a period of 7 days. For treatment of acne, the therapy should be extended for a longer period according to the nature of this disease.

Route of Administration:

For external (Topical) use only

Contraindications:

Hypersensitivity to the preparation.

Infection caused by non-susceptible organisms, in particular, *Pseudomonas aeruginosa*.

Warning and Precautions:

Bacterial resistance has been reported to occur with the use of fusidic acid applied topically. As with all topical antibiotics, extended or recurrent applications may increase the risk of contact sensitisation and the development of antibiotic resistance.

Fusidic acid does not appear to cause conjunctival irritation in experimental animals. Caution should still be exercised, however, when fusidic acid is used near the eyes.

Drug interaction:

None known

Pregnancy and Lactation:

There is inadequate evidence of safety in human pregnancy. Animal studies and many years of clinical experience have suggested that fusidic acid is devoid of teratogenic effect. There is evidence to suggest that when given systemically, fusidic acid can penetrate the placental barrier. The use of topical Fucid in pregnancy requires that the potential benefits be weighed against the possible hazards to the foetus.

Safety in nursing mothers has not been established. When fusidic acid (as the sodium salt) has been given systemically, levels have been detected in breast milk, but with topical use the possible amount of drug present is unlikely to affect the infant.

Side Effects:

Hypersensitivity reactions to the active ingredient in the form of skin rashes; mild stinging and irritation on application have been reported rarely.

Overdosage:

Not applicable

Effects on Ability to Drive and Use Machine:

Not applicable

Storage:

Store below 30°C.

For external use only

Keep container tightly closed.

Keep out of reach of children.

Jauhkan daripada capaian kanak-kanak

Presentation:

Aluminium tubes of 5 g, 10 g, & 15 g.

Manufactured by:

Noripharma Sdn. Bhd. 200701034604 (792633-A)

Lot 5030, Jalan Teratai,

5 ½ Mile Off Jalan Meru,

41050 Klang, Selangor, Malaysia

Date of revision: January 2024