

A combination treatment is recommended (i.e vaginal cream and vaginal tablet) if the labia and adjacent areas are simultaneously infected.

The sexual partner should also undergo local treatment using an external cream if symptoms, e.g. pruritus, inflammation, etc. are present.

Treatment during the menstrual period should not be performed. The treatment should be finished before the onset of menstruation.

Do not use tampons, intravaginal douches, spermicides or other vaginal products while using this product.

Avoidance of vaginal intercourse is recommended in case of vaginal infection and while using this product because the partner could become infected and the effectiveness and safety of latex products such as condoms and diaphragms may be reduced.

During pregnancy, the vaginal tablets should be used and inserted without using an applicator. Intended for use by adults and children 12 years of age and older.

Overdosage

No risk of acute intoxication is seen as it is unlikely to occur following a single vaginal application of an overdose (application over a large area under conditions favorable to absorption) or inadvertent oral ingestion. There is no specific antidote.

If you forget to use

When you remember, continue with your course of treatment. If you have forgotten for more than one day, your infection may not be fully treated, so if symptoms persist upon completion of this course, see your doctor.

Effects on Ability to Drive and Use Machines

The medication has no or negligible influence on the ability to drive or use machinery.

UNDESIRABLE EFFECTS

Frequency not known. The following adverse reactions have been identified during post-approval use of Clotrimazole. As these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency.

- Immune system disorders: anaphylactic reaction, angioedema, hypersensitivity
- Vascular disorder: syncope, hypotension
- Respiratory, thoracic and mediastinal disorders: dyspnea
- Reproductive system and breast disorder disorders: vaginal exfoliation, vaginal discharge, vulvovaginal pruritus, vulvovaginal erythema, vulvovaginal discomfort, vulvovaginal burning sensation, vulvovaginal pain, vaginal haemorrhage
- Gastrointestinal disorders: abdominal pain, nausea
- Skin and Subcutaneous Tissue Disorders: rash, urticaria
- General disorders and administration site conditions: application site irritation, oedema, pain

STORAGE CONDITION

Do not store above 30°C.

Pharmacological Properties

ATC Code: G01A F02

Clotrimazole is an imidazole derivative with a broad spectrum antimycotic activity.

Mechanism of Action

Azoles (e.g. clotrimazole) are usually recommended for the local treatment of vulvovaginal candidosis that is characterized by vulvovaginal symptoms such as itching, burning, discharge, redness, swelling and soreness. Clotrimazole acts against fungi by inhibiting ergosterol synthesis. Inhibition of ergosterol synthesis leads to structural and functional impairment of the cytoplasmic membrane.

Pharmacodynamics Properties

Clotrimazole has a broad antimycotic spectrum of action in vitro and in vivo, which includes dermatophytes, yeasts, moulds, etc.

Under appropriate test conditions, the MIC values for these types of fungi are in the region of less than 0.062-8.0 g/ml substrate.

The mode of action of clotrimazole is primarily fungistatic or fungicidal depending on the concentration of clotrimazole at the site of infection. In-vitro activity is limited to proliferating fungal elements; fungal spores are only slightly sensitive. In addition to its antimycotic action, clotrimazole also acts on gram-positive microorganisms (Streptococci/ Staphylococci), and gram-negative microorganisms (Bacteroides/Gardnerella vaginalis). In vitro clotrimazole inhibits the multiplication of Enterococci – in concentrations of 0.5-10 g/ml substrate and exerts a trichomonacidal action at 100 g/ml. Primarily resistant variants of sensitive fungal species are very rare; the development of secondary resistance by sensitive fungi has so far only been observed in very isolated cases under therapeutic conditions.

Pharmacokinetic Properties

Pharmacokinetic investigations after vaginal application have shown that only a small amount of clotrimazole (3-10%) is absorbed. Due to the rapid hepatic metabolism of absorbed clotrimazole into pharmacologically inactive metabolites the resulting peak plasma concentrations of clotrimazole after vaginal application of a 500mg dose were less than 10ng/ml, suggesting that clotrimazole applied intravaginally is unlikely to lead to measurable systemic effects or side effects.

Incompatibilities

None known

Presentation

Pack of 1 vaginal tablet with applicator.

Manufactured by:

Haupt Pharma Wülfing GmbH
Bethelner Landstrasse 18
31028 Gronau/Leine Germany

Product Registration Holder:

Bayer Co. (Malaysia) Sdn Bhd
25-03 & 25-04, Level 25, IMAZIUM, No. 8, Jalan SS21/37,
Damansara Uptown, 47400 Petaling Jaya, Selangor.

Date of Revision: 01 June 2022

Bayer

AW.CSA/MUP-54450-01

BAYER GP Grenzach Produktions GmbH		Contract manufact.		GP Grenzach		Used template		LFT_200x315mm_Template_CA-21		No. of proofs	
typo pharma Typopharma GmbH Mühlstr. 57 D-79539 Lörrach www.typopharma.com		Contr. manufact. no.		N/A		Min. fontsize text		11 pt		1 23.06.2022 ms 8	
		Product		PIL, CANESTEN VT1 APP MY		Font used		Verlag (Book, Bold)		2 30.06.2022 ms 9	
		Material no.		AW.CSA/MUP-54450-01		Printing colours		Black		3 10	
		Replaced no.		N/A						4 11	
		Order file		N/A						5 12	
typopharma no: # 147641		Country		MY						6 13	
		Format/dimensions		200 x 315 mm		Technical colours		Cutting		7 14	
		Page no.		Page 2 of 2							
		Drawing no.		D0011104							
		Code no. SEM		FPO		N/A					