

MUCORID-P SYRUP

Composition:

Each 5ml contains Carbocisteine 100mg, Promethazine Hydrochloride 2.5mg and Methylparaben (as preservative) 7.5mg.

Description:

A brown colored syrup having fruity flavor.

Pharmacodynamic:

Carbocisteine is a mucolytic agent which decreases the pathological viscosity of the mucus, fluidifies it and facilitates its evacuation, thus promoting recovery. As a mucoregulatory agent, it helps generate the impaired mucosa.

Promethazine is a long acting antihistamine with a mild atropine-like anticholinergic effect and because of its marked effect on the central nervous system, is of great value as an antiemetic, hypnotic, tranquilizer, a potentiator of anaesthetic, hypnotics, sedatives and analgesics.

Pharmacokinetic:

Carbocisteine is rapidly and well absorbed from the gastrointestinal tract with peak plasma concentrations occurring 90 to 120 minutes after an oral dose. It appears to penetrate into lung tissue and respiratory mucus.

Carbocisteine is excreted in the urine as unchanged drug and metabolites. Acetylation, decarboxylation and sulfoxidation have been identified as the major metabolic pathways.

Sulfoxidation may be governed by genetic polymorphism.

Promethazine is well absorbed from the gastrointestinal tract. Clinical effects are apparent within 20 minutes after oral administration and generally last four to six hours, although they may persist as long as 12 hours. Promethazine is metabolized by the liver to a variety of compounds; the sulfoxides of promethazine and N-demethylpromethazine are the predominant metabolites appearing in the urine.

Indication:

Symptomatic treatment of troublesome, non-productive cough, particularly occurring at night.

Recommended dosage:

FOR ADULTS AND CHILDREN OVER THE AGE OF 2 YEAR ONLY.

Symptomatic treatment should be for a short period (a few days) and restricted when cough occurs.

Adults: 3 to 4 tablespoons (1 tablespoon = 15ml) per day, in divided doses.

Children:

- 25 to 30 months: 3 to 4 of the 5ml measuring-spoonfuls per day, in divided doses.
- 30 months to 12 years: 4 to 6 of the 5ml measuring-spoonfuls per day, in divided doses.
- 12 to 15 years: 6 to 9 of the 5ml measuring spoonfuls per day, in divided doses.

This medicine should preferably be taken in the evening, because of the marked sedative effect of promethazine.

Route of Administration:

Oral administration

Contraindication:

MUCORID-P is contraindicated for the following:

- Hypersensitivity to any of the excipients especially methylparaben
- History of agranulocytosis
- Risk of urine retention related to urethral or prostate disorders
- Risk of angle-closure glaucoma

Concomitant use of sultopride is generally inadvisable.

Contraindicated in children below 2 years of age.

Warning and precaution:

Caution is recommended in patients suffering from gastroduodenal ulcers.

As phenothiazines have been considered as hypothetical risk factors in the onset of sudden death in infant, their use is not recommended in infants with a risk of apnea below the age of 1 year.

Epileptic patients should be monitored because of a possible decrease of the epileptogenic threshold.

If used in children, the possibility of bronchial asthma or gastroesophageal reflux should be ruled out before using promethazine as an antitussive.

As phenothiazines have photosensitizing effect, patients should try not to expose themselves to the sun during treatment.

Promethazine hydrochloride should be used with care:

1. In the elderly

- Increased sensitivity to orthostatic hypotension, dizziness and sedation.
- Chronic constipation (risk of paralytic ileus)
- Possible prostate hypertrophy

2. In patients with certain cardiovascular disorders, because of the tachycardiac and hypotensive effect of phenothiazines

3. In severe liver and/or kidney insufficiency (because of the risk of accumulation)

This preparation contains Sodium metabisulphite that may cause serious allergic type reactions in certain susceptible patients.

Do not use if known to be hypersensitive to bisulphites.

This product contains Promethazine Hydrochloride. It should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression.

Warning: Not suitable to be used in children below 2 years old.

For children between 2 to 6 years old – use with caution and follow the advice given by the doctor or pharmacist.

Drug interaction:

- Alcohol

Alcohol enhances the sedative effect of antihistamine. Impaired alertness can make driving or machine operation dangerous.

- Sultopride

Increase risk of ventricular rhythm disorders, particularly torsades de pointes, by addition of electrophysiological effects.

- Other central nervous system depressants
Sedative antidepressants, barbiturates, clonidine and related substances, hypnotics, morphine derivatives (analgesics and antitussives), methadone, neuroleptics, anxiolytics. Enhancement of central depression. Impaired alertness can make driving or machine operation dangerous.
- Atropine and other atropine substances
Imipramine antidepressants, anticholinergic antiparkinsonians, atropine antispasmodics, disopyramide, phenothiazines neuroleptics; addition of adverse atropine effects such as urine retention, constipation, dry mouth.

Pregnancy and lactation:

MUCORID-P should be avoided during the first trimester as a precautionary measure. Then, it should only be prescribed if necessary. During the last trimester of pregnancy it should only be used occasionally.

If MUCORID-P is administered at the end of pregnancy, it may be justified to ensure a period of monitoring of the neurological and gastrointestinal functions of the neonate. There is a low but detectable excretion of antihistamines into breast milk. Therefore, the use of this medicine should be avoided in the event of breast-feeding.

Side effects:

Gastrointestinal intolerance symptoms (gastric pain, nausea, diarrhoea) have been reported. To overcome this problem, dosage reduction is recommended.

The following is related to the presence of promethazine:

- Sedation or drowsiness, usually during the initial treatment
- Anticholinergic effects for instance dryness of mucosa, constipation, accommodation disorders, mydriasis, cardiac palpitation, risk of urine retention
- Orthostatic hypotension
- Equilibrium disorders, dizziness, impaired memory or concentration
- Motor in-coordination, tremor (more frequent in elderly patients)
- Mental confusion, hallucination
- Rarely in infants, excitation effects: agitation, nervousness, insomnia
- Erythema, eczema, pruritus, purpura and urticaria
- Edema, more rarely Quincke's oedema
- Anaphylactic shock
- Photosensitization
- Leucopenia, neutropenia and exceptional cases of agranulocytosis
- Thrombocytopenia
- Haemolytic anemia

Immune System Disorders:

Anaphylactic / anaphylactoid reaction

Skin and Subcutaneous Tissue Disorders:

Severe cutaneous adverse reactions (SCAR) e.g. erythema multiforme, Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN). In most of these cases reported at least one other drug was administered at the same time, which may have possibly enhanced the described mucocutaneous effects.

Overdosage:*Carbocisteine:*

Gastrointestinal disturbance is the most likely symptom of overdose.

If overdose happens, gastric lavage may be beneficial, followed by observation.

Promethazine:

Convulsions (particularly in infants and children), consciousness disorders, coma.

Symptomatic treatment should be initiated in a specialized unit.

Effects on ability to drive and use machines:

Attention is drawn especially for those who drive or use a machine, to the risk of drowsiness related to the use of this medicinal product, especially at the start of treatment.

Storage condition:

Store in a dry place below 30°C.

Keep container tightly closed.

Protect from light.

Presentation:

HDPE Bottles of 60ml, 90ml and 120ml.

Manufactured by:

NORIPHARMA SDN. BHD. 200701034604 (792633-A)

Lot 5030, Jalan Teratai,

5 1/2 Mile Off Jalan Meru, 41050

Klang, Selangor Darul Ehsan

Date of revision:

December 2023