

Package Insert

NERVON TABLET

Composition:

Each tablet contains:

Thiamine Mononitrate (Vitamin B ₁)	100mg
Pyridoxine HCl (Vitamin B ₆)	200mg
Cyanocobalamin (Vitamin B ₁₂)	200mcg

Description:

A pink color round shaped film coated tablet with 11mm diameter.

Pharmacology:

B1, B6 and B12 are important vitamins essential for the proper functioning of the nervous system. They are required for the normal metabolic processes of nerve cells. Nervon, as the logical and scientific treatment of nerve degeneration, its associated symptoms and as an adjuvant with NSAIDs to potentiate their analgesic effect with reduced treatment and lesser side effects.

Pharmacokinetic:

Thiamine, Pyridoxine and Cyanocobalamin are absorbed from gastrointestinal tract.

Thiamine is widely distributed to most body tissue; it is not stored in the body. Thiamine is excreted in urine as unchanged form or as its metabolites, pyrimidine.

Pyridoxine is converted to its active form pyridoxal phosphate and transformed to pyridoxic acid which is excreted in the urine.

Cyanocobalamin is bound to plasma proteins and stored in the liver. It is excreted in the bile and undergoes some enterohepatic recycling. It is also excreted in urine.

Indication:

Nervon is indicated for neurological and other disorders associated with disturbance of the metabolic functions influenced by B complex vitamins, including

- diabetic polyneuropathy
- alcoholic peripheral neuritis
- polyneuropathy of pregnancy
- toxemia of pregnancy
- nausea and vomiting of pregnancy
- post-influenza neuropathies

Nervon is also recommended for the treatment of neuritis and neuralgia of the spinal nerves, especially facial paresis, cervical syndrome, low back pain, ischialgia, etc.

Contraindication:

Known hypersensitivity to vitamins B₁, B₆ or B₁₂.

Should not be used in patients on Levodopa therapy.

Dosage and Directions:

Take 1 tablet 2-3 times daily to treat moderate cases. The coated tablets are swallowed unchewed with liquid or after meals.

Warning & Precaution:

Cyanocobalamin should not be given before a diagnosis has been fully established because of the possibility of masking symptoms of subacute degeneration of the spinal cord.

Cyanocobalamin is not a suitable form of vitamin B₁₂ for the treatment of optic neuropathies associated with raised plasma concentrations of cyanocobalamin.

Long term administration of large doses of pyridoxine is associated with the development of severe peripheral neuropathies. This may occur with doses in excess of about 2g daily. Cyanocobalamin should not be used in megaloblastic anaemia of pregnancy. Administration of doses greater than 10 microgram daily may produce a haematological response in patients with folate deficiency.

Pregnancy and Lactation

If you are pregnant or breastfeeding, please consult your doctor or healthcare professional before taking this product.

Adverse Reaction:

Allergic hypersensitivity reactions have occurred rarely following parenteral administration of vitamin B₁₂ compounds. Adverse effects seldom occur following administration of vitamin B₁, but hypersensitivity reactions have occurred, this is mainly after parenteral administration. Long-term administration of large doses of pyridoxine (in excess of 2g daily) is associated with the development of severe peripheral neuropathies

Interaction:

Loop diuretics (furosemide, ethacrynic acid, bumetanide): Chronic use of loop diuretics may result in thiamin deficiency. Concomitant intake of thiamin with food that contains sulfites may inactivate thiamin. Concomitant intake of thiamin with tea, coffee and decaffeinated coffee may inactivate thiamin. This can be reversed by taking supplements.

Pyridoxine reduces the effects of levodopa, but this does not occur if a dopa decarboxylase inhibitor is also given. Pyridoxine reduces the activity of altretamine. It has also been reported to decreased serum concentrations of Phenobarbital and phenytoin. Many drugs may increase the requirement of pyridoxine; such drugs include hydralazine, isoniazid, penicillamine, and oral contraceptives.

Absorption of vitamin B₁₂ from the gastrointestinal tract may be reduced by neomycin, aminosalicic acid, histamine H₂-antagonists, metformin, para-aminosalicylic acid, potassium chloride, colchicines, and proton pump inhibitors (lansoprazole, omeprazole, pantoprazole, rabeprazole). Cholestyramine and colestipol may decrease the enterohepatic reabsorption of vitamin B₁₂. Inhalation of the anesthetic agent nitrous oxide can produce a functional vitamin B₁₂ deficiency. The use of antibiotics may alter the intestinal microflora and may decrease the possible contribution of vitamin B₁₂ by certain inhabitants of microflora to the body's requirement for the vitamin.

Overdosage:

Treatment is symptomatic and supportive.

Shelf life:

2 years

Storage Condition:

Store in dry place below 30°C.

Presentation:

Blister of 10 capsules. Box of 1, 3, 6, 10 and 50 blisters.

Manufacturer:

Pahang Pharmacy Sdn. Bhd. (10826-K)
Lot 5979, Jalan Teratai 5 ½ Miles Off Jalan Meru,
41050 Klang, Selangor,
Malaysia.

Date revised: 091214