

13 cm.

25 cm.

AMTUSS 30 mg/5 mL Syrup

MUCOLYTIC

Each 5 mL contains:
Ambroxol hydrochloride 30 mg

❖ **PRODUCT DESCRIPTION:**

Clear, colorless solution with grape flavor

❖ **MECHANISM OF ACTION:**

Pharmacology

Preclinically, Ambroxol has been shown to increase respiratory tract secretion. It enhances pulmonary surfactant production and stimulates ciliary activity. These actions result in improved mucus flow and transport (mucociliary clearance). Improvement of mucociliary clearance has been shown in clinical pharmacologic studies. Enhancement of fluid secretion and mucociliary clearance facilitates expectoration and eases cough.

❖ **INDICATION:**

Secretolytic therapy in acute and chronic bronchopulmonary diseases associated with abnormal mucus secretion and impaired mucus transport.

❖ **DOSAGE AND ADMINISTRATION:**

Oral

Adult and Children >12 years:

2 teaspoonfuls 2 times daily.

This regimen is suitable for the initial treatment of chronic conditions up to 14 days.

Children <12 years:

The following dosage regimen is recommended depending on the severity of the disease.

Children 6-12 years : 1 teaspoonful 2-3 times daily

2-6 years: ½ teaspoon 3 times daily

1-2 years: ½ teaspoon 2 times daily

This dosage regimen is for initial treatment; the dosage may be halved after 14 days.

AMTUSS liquid should be taken at meal times.

❖ **SIDE EFFECT:**

AMTUSS is generally well tolerated. Mild upper gastrointestinal side effects (primarily pyrosis, dyspepsia and occasionally nausea, vomiting) have been reported, principally following parenteral administration. Allergic reactions have occurred rarely, primarily skin rashes.

There have been extremely rare case reports of severe acute anaphylactic-type reactions but their relationship to Ambroxol is uncertain. Some of these patients have also shown allergic reactions to other substances.

❖ **CONTRAINDICATION:**

AMTUSS should not be used in patients known to be hypersensitive to Ambroxol or other components of the formulation.

❖ **PREGNANCY AND LACTATION:**

Preclinical studies as well as extensive clinical experience after the 28th week have shown no evidence of ill-effects during pregnancy. Nonetheless, the usual precautions regarding the use of drug during pregnancy, especially during the 1st trimester, should be observed. During lactation, AMTUSS should be used only if absolutely necessary.

❖ **DRUG INTERACTION:**

Administration of Ambroxol together with antibiotics (Amoxicillin, Cefuroxime, Erythromycin, Doxycycline) leads to higher antibiotic concentration in the lung tissue. No clinically relevant unfavorable interaction with other medications have been reported.

❖ **OVERDOSE AND TREATMENT:**

No symptoms of overdosage have been reported in man to date. If they occur, symptomatic treatment should be provided.

❖ **SHELF LIFE:**

Two (2) years

❖ **STORAGE:**

Store at temperature of not more than 30°C.

❖ **SUPPLY:**

Bottle 60 mL

Manufactured by:

UNISON LABORATORIES CO., LTD.

39 Moo 4, Klong Udomcholjorn, Muang Chachoengsao,
Chachoengsao 24000 Thailand

Marketing Authorization Holder/ Importer:

MEDISPEC (M) SDN. BHD.

No. 55 & 57, Lorong Sempadan 2, (Off Boundary Road)
11400 Ayer Itam, Penang, Malaysia

PM Specification	Product Name	Code No.	Dimension	Packaging Type	Thickness
	AMTUSS Syrup	IAML 0011	W 13 x L 25 cm.	กระป๋องชนิด	60 g (0.08 mm.)
Design by:		Checked by:		Approve by:	
				(Only Pattern and Colors)	