

SIMCONE-T SUGAR FREE

CHEWABLE TABLET

CONTENT

Each chewable tablet contains 80.0mg of simethicone.

Excipients: Aspartame, sorbitol, povidone, menthol, peppermint oil & magnesium stearate.

DESCRIPTION

A white, round, biconvex tablet.

Flavouring: Menthol and Peppermint.

PROPERTIES

Pharmacodynamic

Simethicone is liquid dimethicone activated with finely divided silicon dioxide to enhance the defoaming properties of the silicone. Simethicone acts by decreasing the surface tension of gas bubbles, thus facilitating their coalescence and expulsion as flatus or belching. It also prevents the formation and accumulation of mucus-enclosed pockets of gas in digestive tract. Simethicone facilitates the passage of gas through bowel lumen and allows patients to excrete a greater volume of gas at one time, thereby reducing the number of flatus events. Thus, less residual gas is present to cause uncomfortable or painful pressure in the stomach and intestines.

This sugar free formulation product contains aspartame as a sweetener agent and hence, it is suitable for diabetic patients and calories-conscious users.

Pharmacokinetics

Simethicone is pharmacologically inert and is not absorbed from the gastrointestinal tract nor does it interfere with gastric secretion or absorption of nutrients. Following oral administration, the drug is excreted in faeces unchanged.

INDICATION

Simcone-T Sugar Free chewable tablet is indicated for the relief of discomfort and pain caused by excessive gas in the stomach and intestines as in flatulence, dyspepsia, meteorism, and post-operative gaseous distension. **Simcone-T Sugar Free** chewable tablet can be used to eliminate gas, air and foam from the gastrointestinal tract prior to endoscopic examination or radiography.

RECOMMENDED DOSAGE

For oral administration.

Adult and teenager: 80mg (1 tablet) to 160 mg (2 tablets) after each meal and at bedtime. Unless otherwise directed by the doctor, the total dose should not exceed 480 mg (6 tablets) per day.

Children (2 – 12 years): 40 mg (half-tablet) after each meal and at bedtime, up to a maximum of 240 mg (3 tablets) per day.

Chew the tablets thoroughly before swallowing to enable the medicine to work faster and more completely.

Missed dose: Take the missed dose as soon as you remember. However, if it is time for the next dose, skip

the missed dose and only take your next regularly scheduled dose.

CONTRAINDICATION

Simcone-T Sugar Free chewable tablet should not be taken by patients who have previous allergic reaction to simethicone or to any of its excipients.

Simcone-T Sugar Free chewable tablet is also contraindicated in patients with known or suspected intestinal perforation and obstruction.

WARNING / PRECAUTION

Inform the doctor if you have any type of serious illness or allergy, are pregnant or are breast-feeding.

Phenylketonurics are warned that this product contains phenylalanine.

DRUG INTERACTION

As simethicone reduces the bioavailability of phenytoin slightly, take **Simcone-T Sugar Free** chewable tablet 2 hours before or after phenytoin.

PREGNANCY & LACTATION

Simethicone is safe for use during pregnancy and lactation. Nevertheless, phenylketonurics should be warned that this product contains phenylalanine.

SIDE EFFECTS

Although oral administration of simethicone is generally well tolerated when used as directed, the following adverse reactions have been observed on rare occasions:- bloating, constipation, diarrhoea, gas, heartburn.

SYMPTOMS & TREATMENT OF OVERDOSE

As simethicone is not absorbed into the body, there is no likelihood of overdosage.

STORAGE CONDITIONS

Do not store above 30°C.

Protect from heat and light.

Keep out of the reach of children.

Jauhi daripada kanak-kanak.

Shelf life: Please refer to outer box label.

PACKING / PACK SIZES

Box of (10 x 10) and (50 x 10) tablets in aluminium foil strip pack.

Manufactured by:

T.O. Pharma Co., Ltd.

101, Ladprao 124,

Bangkok 10310, Thailand.