

DISUF CREAM

VIDIS11-1 (MY)

Description

White, opaque cream.

Composition

Fusidic Acid 2.0% w/w.

Actions And Pharmacology

Fusidic acid is an antibiotic with a bacteriostatic or bactericidal activity, mainly against Gram positive bacteria.

Fusidic acid inhibits bacterial protein synthesis although, in contrast to drugs such as the macrolides or tetracyclines, it does not bind to the bacterial ribosomes, but inhibits a factor necessary for translocation of peptide subunits and elongation of the peptide chain.

Pharmacokinetics

In vitro studies show that fusidic acid can penetrate intact human skin. The degree of penetration depends on factors such as the duration of exposure to fusidic acid and the condition of the skin. Fusidic acid is excreted mainly in the bile with little excreted in the urine.

Indications

Disuf Cream is indicated for the treatment of skin infections caused by *Staphylococci*, *Streptococci*, *Propionibacterium acnes*, *Corynebacterium minutissimum* and other organisms sensitive to fusidic acid. The most important indication being: Impetigo, Boils, Carbuncles, Paronychia, Infected wounds, Sycosis barbae, Hidradenitis, Erythrasma, Folliculitis.

Contraindications

- Infections caused by non-susceptible organisms.
- Patients with hypersensitivity to any of the ingredients.

Warning / Precautions

- Caution should be observed when applying Disuf Cream in the eye region as this preparation may cause irritation if it gets into the eye.
- Bacterial resistance has been reported to occur with the use of fusidic acid applied topically.
- As with all topical antibiotics, extended or recurrent application may increase the risk of contact sensitisation and the development of antibiotic resistance.

Pregnancy And Lactation

There is inadequate evidence of safety in human pregnancy. Safety in nursing mothers has not been established.

Main Side / Adverse Effects

Hypersensitivity reactions to the active ingredient in the form of skin rashes; mild stinging and irritation on application have been reported rarely.

Overdose

Not applicable.

Dosage And Administration

Disuf Cream is applied to the affected area 2-3 times daily, generally for a period of 7 days.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage : Store below 30°C.
Presentation/Packing : 15 g aluminium tube.

Product Registration Holder /
Manufactured by: HOVID Bhd.
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Revision date: January 2017

hovid