

Simcone Suspension 67 mg/ml

CONTENT

Each 1ml contains simethicone 67.0mg.

Excipients: Sorbitol, microcrystalline cellulose, carboxymethylcellulose sodium, sodium citrate, citric acid, bronopol, myacide SP & propylene glycol.

DESCRIPTION

White aqueous suspension.

PROPERTIES

Pharmacodynamic

Simethicone is liquid dimethicone activated with finely divided silicon dioxide to enhance the defoaming properties of the silicone. Simethicone acts by decreasing the surface tension of gas bubbles, thus facilitating their coalescence and expulsion as flatus or belching. It also prevents the formation and accumulation of mucus-enclosed pockets of gas in digestive tract. Simethicone facilitates the passage of gas through bowel lumen and allows patients to excrete a greater volume of gas at one time, thereby reducing the number of flatus events. Thus, less residual gas is present to cause uncomfortable or painful pressure in the stomach and intestines.

Pharmacokinetics

Simethicone is pharmacologically inert and is not absorbed from the gastrointestinal tract nor does it interfere with gastric secretion or absorption of nutrients. Following oral administration, the drug is excreted in faeces unchanged.

INDICATION

Simcone Suspension is indicated for the relief of discomfort and pain caused by excessive gas in the stomach and intestines as in flatulence, dyspepsia, meteorism, and post-operative gaseous distension. **Simcone Suspension** can be used to eliminate gas, air and foam from the gastrointestinal tract prior to endoscopic examination or radiography.

RECOMMENDED DOSAGE

For oral administration.

Adults and teenagers: 0.60ml – 1.40ml, four times daily with or after each meal. The dose should not be more than 480mg/24 hours.

Infants: 0.25ml – 0.50ml, three times daily with or after each meal. The dose should not be more than 240mg/24 hours. **Simcone Suspension** may be added to baby's formula or given directly from a dropper.

Shake well before use.

CONTRAINDICATION

Simcone Suspension should not be taken by patients who have previous allergic reaction to simethicone or to any of its excipients.

Simcone Suspension is also contraindicated in patients with known or suspected intestinal perforation and obstruction.

WARNING / PRECAUTION

Inform the doctor if you have any type of serious illness or allergy, are taking other medicines, pregnant or are breast-feeding.

DRUG INTERACTION

As simethicone reduces the bioavailability of phenytoin slightly, take **Simcone Suspension** two hours before or after phenytoin.

PREGNANCY & LACTATION

Simethicone is safe for use during pregnancy and lactation.

SIDE EFFECTS

Although oral administration of simethicone is generally well tolerated when used as directed, the following adverse reactions have been observed on rare occasions: - bloating, constipation, diarrhoea, gas, heartburn.

SYMPTOMS & TREATMENT OF OVERDOSE

As simethicone is not absorbed into the body, there is no likelihood of overdosage.

STORAGE CONDITIONS

Do not store above 30°C.

Keep in tight containers.

Shake well before use.

Keep out of the reach of children.

Jauhi daripada kanak-kanak.

Shelf life: Please refer to outer box label.

PACKING / PACK SIZES

Bottle of 15ml.

Box of 12 bottles of 15ml.

Manufactured by:

T.O. Pharma Co., Ltd.

101 Soi Ladprao 124 (Sawatdikan),

Ladprao Road, Phlapphla, Wangthonglang,

Bangkok 10310 THAILAND.

Revision: June 2025.