



Betex SC Tablet

Name of Product

Betex SC Tablet

Strength of the medicine

Each tablet contains :

Thiamine Disulfide (Vitamin B1)	100 mg
Pyridoxine HCl (Vitamin B6)	200 mg
Cyanocobalamin (Vitamin B12)	200 mcg

Description

Round, yellow, biconvex, sugar-coated tablets, with diameter of 12.7 mm and thickness of 6.7 mm, no superficial marking.

Pharmacology

Thiamine is involved in carbohydrate metabolism and in nerve transmission. Pyridoxine is involved in amino acid metabolism and in protein metabolism. Cyanocobalamin influences hematopoiesis. It is involved in formation and metabolism of purines and pyrimidines and in the synthesis of nucleoproteins.

Pharmacokinetic properties

Thiamine is absorbed from the gastrointestinal tract and is widely distributed to most body tissues. It is not stored in the body and amounts in excess of the body's requirements are excreted in urine as unchanged thiamine or as its metabolite, pyrimidine.

Pyridoxine is absorbed from the gastrointestinal tract and is converted to the active form, pyridoxal phosphate. Pyridoxal phosphate is transformed to pyridoxic acid and excreted in the urine. Cyanocobalamin is absorbed from the gastrointestinal tract. In the blood, it is bound to plasma proteins and stored in the liver. It is excreted in the bile and undergoes some enterohepatic recycling; also excreted in urine.

Indications

- Vitamin B1, B6, B12 deficiency, neuritis, and other disorders associated with vitamin B1, B6, B12
- Neural Disturbance: Alcoholic neuritis, diabetic neuritis. Peripheral neuritis.

Recommended Dosage

1 tablet 3 times daily after meals or as directed by the physician.

Mode of administration

Oral

Contraindications

Contraindicated for use in cases with known hypersensitivity to any of the ingredients.

Precautions

Cyanocobalamin is not a suitable form of Vitamin B12 for the treatment of optic neuropathies associated with raised plasma concentration of cyanocobalamin.

Interactions

Pyridoxine decreases the action of levodopa.

Use in pregnancy and lactation

Pregnancy : The possibility of fetal harm appears remote. Use during pregnancy only if clearly needed.

Lactation : Use with caution in nursing women.

Adverse Reactions

Pruritus, urticaria, weakness, sweating, nausea, restlessness.

Overdose and Treatment

Symptoms of overdose is very rare, treatment is symptomatic and supportive.

Overdosage of Pyridoxine HCl

Patient receiving 2 to 7 g/day (or > 0.2 g/day for > 2 months) have developed sensory neuropathy with associated ataxia and numbness of the hands and feet. When pyridoxine is discontinued, symptoms will lessen. It may take 6 months for sensation to normalize.

Storage Conditions

Store below 30°C and protect from light.

Dosage Forms and packaging available

Tablets ; Boxes of 25 x 10's and 100 x 10's

Manufacturer

The United Drug (1996) Co., Ltd.

208 Romklao Road, Minburi

Bangkok 10510, Thailand

Under the license from FAVOREX Ltd., Switzerland

Diimport dan diedarkan oleh :

Milrin Pharmaceutical Co. (M) Sdn. Bhd.

125-L, 5 th Floor, Wisma Getah,

Macalister Road, 10400

Penang, Malaysia.

Fax : 604-2286042, 2286741 & 2288644

Fax : 604-2280757

The date of revision : October 2017