

Revision Dt. 25-02-2023
Revision Dt. 05-08-2024

Artwork is in same size
Size: 90 mm x 188 mm

For Hospital or Medical practitioner or Laboratory

HEXIGEL[®] 1% w/w

(Chlorhexidine gluconate 1.0% w/w)

Name and strength of Active ingredient(s):

Chlorhexidine gluconate solution BP 5% v/w equivalent to chlorhexidine gluconate 1.0% w/w

Description:

A clear colourless transparent gel with characteristic pleasant odour and taste.

Indication

For oral hygiene, relief of gingivitis, mouth ulcers and oral infections.

Recommended Dose

To be brushed on the teeth once or twice daily for oral hygiene, plaque inhibition and gingivitis. To be applied on the affected areas, once or twice daily, for management of aphthous and other oral ulcers.

Contraindication

Hexigel is contraindicated for patients who have previously shown a hypersensitivity reaction to chlorhexidine. However, such reactions are extremely rare.

Warning and precaution

Brown staining may be reported with extended use in some cases, which may be removed by brushing or polishing. NOT TO BE SWALLOWED.

Avoid contact with eyes- if Chlorhexidine solutions come into contact with eyes, wash out promptly and thoroughly with water. As toothpastes may contain anionic surfactants such as sodium laurilsulfate, which are incompatible with chlorhexidine, it has been recommended that at least 30 minutes should be allowed to elapse between the use of toothpaste and oral chlorhexidine preparations.

Hexigel contains chlorhexidine. Chlorhexidine is known to induce hypersensitivity, including generalized allergic reactions and anaphylactic shock. The prevalence of chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare. Hexigel should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine.

If any signs or symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness and rapid heart rate develop, immediately stop using the product. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Interaction with other medicaments

Chlorhexidine is incompatible with anionic agents.

Undesirable Effects / Side effects

Immune system disorders : Frequency not known : Hypersensitivity including anaphylactic shock. A superficial discoloration of the dorsum of the tongue may occur. This disappears after treatment is discontinued. Discoloration of the teeth and silicate or composite restorations may also occur. The stain is not permanent and can largely be prevented by brushing with conventional toothpaste daily before using the gel. However, in certain cases of professional prophylaxis (scaling and polishing) may be required to remove the stain completely. Stained anterior tooth coloured restorations with poor margins or rough surfaces, which are not adequately cleaned by professional prophylaxis, may require replacement.

Transient disturbance of taste sensation and a burning sensation of the tongue may occur on initial use of the gel. These effects usually diminish with continued use.

Symptoms and treatment of overdose

This has not been reported. Accidental ingestion: Chlorhexidine taken orally is poorly absorbed. Systemic effects are unlikely even if large amounts are ingested. However, gastric lavage may be advisable using milk, raw egg, gelatin or mild soap. Employ supportive measures as appropriate.

Storage condition

Store below 30°C in a dry place. Protect from light. Do not freeze. KEEP TUBE CAPPED WHEN NOT IN USE.

Shelf life

36 months from the date of manufacture

Presentation

7.5/15g multicoloured laminated tube with white, tightly fitted standup cap, Collating and wrapping of 4x3 packed carton 7.5/15g in 12's BOPP pack.

Manufactured By

ICPA Health Products Limited
286/287/288, GIDC, Ankleshwar,
Gujarat 393002,
India.

Revision date: 05/08/24

MAL Number : **MAL08042440XZ**

AP40515R3