

DISUF-H CREAM

VIDIS04-PC3 (MY)

DESCRIPTION

White, opaque cream.

COMPOSITION

Fusidic Acid (Micronised)	2.0% w/w
Hydrocortisone Acetate (Micronised)	1.0% w/w

PHARMACODYNAMICS

Fusidic acid is an antibiotic with a bacteriostatic or bactericidal activity, mainly against Gram positive bacteria. Fusidic acid inhibits bacterial protein synthesis although, in contrast to drugs such as the macrolides or tetracyclines, it does not bind to the bacterial ribosomes, but inhibits a factor necessary for translocation of peptide subunits and elongation of the peptide chain.

Hydrocortisone acetate diffuses across the cell membrane and stimulates transcription of messenger RNA (mRNA) and subsequent protein synthesis of various inhibitory enzymes responsible for the anti-inflammatory effect of topical corticosteroids. These anti-inflammatory effects includes inhibition of early processes such as edema, fibrin deposition, capillary dilatation, movement of phagocytes into the area, and phagocytic activities.

When applied topically, this product is effective against *staphylococci*, *streptococci*, *corynebacteria*, *Neisseria* and certain *Clostridia* and *Bacteroides*.

PHARMACOKINETICS

In vitro studies show that fusidic acid can penetrate intact on human skin. The degree of penetration depends on factors such as the duration of exposure to fusidic acid and the condition of the skin. Hydrocortisone acetate can be absorbed when given locally.

INDICATION

Disuf-H Cream is indicated for the treatment of dermatitis, including atopic dermatitis and contact dermatitis, where an infection with bacteria sensitive to fusidic acid is suspected or confirmed.

CONTRAINDICATIONS

Hypersensitivity to fusidic acid and its salts. As with other topical corticosteroid preparations, Disuf-H Cream is contraindicated in primary bacterial, viral and fungal skin infections, skin manifestations in relation to tuberculosis or syphilis, perioral dermatitis and rosacea.

WARNINGS AND PRECAUTIONS

- Caution in pregnant patients and in patients with tuberculosis of the skin.
- Caution should be exercised when the product is used near the eye, since hydrocortisone acetate can cause glaucoma if the preparation gets into the eye.
- Pediatric therapy continuing for longer than 2 weeks should be evaluated carefully by the physician.
- Long term continuous therapy should be avoided in infants as adrenal suppression may occur.
- Steroid antibiotic combination should not be used for more than 7 days in the absence of any clinical improvement.
- Bacterial resistance has been reported to occur with the use of fusidic acid applied topically.
- As with all topical antibiotics, extended or recurrent application may increase the risk of contact sensitisation and the development of antibiotic resistance.
- As Disuf-H cream contains a corticosteroid it is not recommended in the following conditions: atrophic skin, cutaneous ulcer, acne vulgaris, fragile skin veins and perianal and genital pruritus.
- Contact with open wounds and mucous membranes should be avoided.
- As with all corticosteroids, prolonged use on the face should be avoided.

PREGNANCY AND LACTATION

Topical corticosteroids should not be used extensively, in large amounts or for prolonged periods in pregnant patients, as fetal abnormalities have been seen in animals due to systemic absorption. There is inadequate evidence of safety in human pregnancy. Safety in nursing mothers has not been established. Topical corticosteroids should not be applied to the breast prior to nursing.

DRUG INTERACTION

None known

MAIN SIDE / ADVERSE EFFECTS

Topical use of steroids may result in skin atrophy, telangiectasia or skin striae, especially during prolonged application. As with all other corticosteroids, folliculitis, hypertrichosis, perioral dermatitis, contact dermatitis, depigmentation and systemic activity, although rare, may occur. Mild stinging, irritation and hypersensitivity reactions to the active ingredient in the form of skin rashes have rarely been reported.

OVERDOSE

Acute overdosage is very unlikely to occur. However, chronic overdosage or misuse may result in increased risk of topical or systemic side effects.

DOSAGE AND ADMINISTRATION

For topical use only.

Route of administration
is added.

DISUF-H CREAM

Apply sparingly to the affected area:

- **Uncovered lesions**
2-3 times daily.
- **Covered lesions**
Less frequent application may be adequate.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage:

Store below 30°C.

Presentation/Packing:
15 g aluminium tube.

Product Registration Holder /
Manufactured by: HOVID Bhd.
121, Jalan Tunku Abdul Rahman
(Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Revision date: December 2017

To amend storage condition to
comply with ASEAN
guideline