

Besonin Aqua Nasal Spray

(Budesonide 100mcg/dose)

Besonin Aqua Nasal Spray

(Budesonide 2 mg/mL)

Composition

Budesonide 2 mg/mL: Each mL contains Budesonide 2.0 mg. Each metered dose delivers Budesonide 100 mcg.

Description

Besonin Aqua Nasal Spray is a metered-dose, manual-pump spray formulation containing a micronized suspension of budesonide in an aqueous medium.

It appears as a white to almost white suspension.

Pharmacodynamics

Budesonide is a non-halogenated corticosteroid, which has a potent glucocorticoid activity and weak mineralocorticoid activity. As a measure of systemic activity, budesonide is 40 times more potent than cortisol when administered subcutaneously and 25 times more potent when administered orally in the rat thymus involution assay. The precise mechanism of corticosteroid actions in seasonal and perennial allergic and nonallergic rhinitis is not known. Corticosteroids have been shown to have a wide range of inhibitory activities against multiple cell types and mediators involved in allergic- and non-allergic/irritant-mediated inflammation. These anti-inflammatory actions of corticosteroids may contribute to their efficacy in asthma. Budesonide has potent local anti-inflammatory effects on respiratory mucosa when administered topically. It works by decreasing inflammation (swelling) in the nasal passages caused by allergens.

Pharmacokinetics

Bioavailability of oral budesonide in man is low due to an extensive first pass metabolism in the liver.

The systemic availability of budesonide from Besonin Aqua, with reference to the metered dose is 33%. AUC after administration of 256 micrograms budesonide from Besonin Aqua is 2.7 nmol/L in adults.

Indication

Prevention and treatment of seasonal and perennial allergic rhinitis and vasomotor rhinitis.

Dosage

Dose for Adults and Children > 6 Years.

100 mcg nasal spray: One spray into each nostril daily.

Directions for use

1. This medicine is for intranasal use only. Shake the bottle gently before use.
2. Prime for initial use by putting two fingers on the shoulders of the spray pump unit and place your thumb on the bottom of the bottle. Press upward with thumb, release, and repeat until a fine mist appears (5-10 sprays). Now your pump is primed and ready to use. If 2 or more days have elapsed since the last use, the pump should be reprimed with 2 sprays or until a fine mist appears.
3. Gently blow nose to clear nostrils.
4. Place the spray tip 0.5 cm to 1 cm into one nostril. Hold bottle vertically upright, allowing spray tip to aim toward the back of the nose. Close other nostril with finger, rapidly press once with thumb and sniff gently at the same time.

5. Repeat in the other nostril.
6. Do NOT take more than the recommended dose of this medicine.
7. Patients should use medicine at regular intervals since its effectiveness depends on its regular use.
8. Missed dose: Take your next dose as soon as you remember. If it is time for your next dose, skip the missed dose and go back to your regular schedule. Do not double doses.

Precaution

1. Avoid getting this medicine in your eyes.
2. Anyone who has chicken pox or measles should not take this medicine.
3. Intranasal corticosteroids may affect growth rates in children. Talk to your doctor for more information.

Pediatric: After administration of budesonide, the time to reach peak concentrations and plasma half-life were similar in children and in adults. Children had plasma concentrations approximately twice those observed in adults due primarily to differences in weight between children and adults.

Pregnancy/Nursing: If you are planning to become pregnant or want to breast-feed, talk to your doctor first to discuss the benefits and risks associated with this medicine. For breast-feeding mothers, it is NOT known if this medicine is excreted in the breast milk. Do NOT use if you are nursing until you have talked with your doctor, or pediatrician.

Overdosage:

Acute overdose should not present clinical problems.

Contraindication

Hypersensitivity to any of the ingredients in these preparations contraindicates its use.

Drug interaction

Let your doctor know if you are taking any other

medications either by prescription or over-the-counter. There are no serious documented drug interactions with this medicine.

Side effect

Side effects caused by this medicine include: nasal irritation, nasal dryness, epistaxis, coughing, sore throat, hoarseness, bad taste, and nausea.

Storage

100 mcg/dose: Store below 30°C.

Package

Besonin Aqueous Nasal Spray 100mcg/dose is supplied in a 15 ml HDPE plastic bottle, pump and nasal applicator. Pack size of 200 sprays.

Reg. No.:

100 mcg/dose MAL08082702AZ

Manufactured by

**SYNMOSA BIOPHARMA CORPORATION,
Synmosa Plant**

No.6, Kuang Yeh 1st Road, Hu-Kuo Hsiang,
Hsin-Chu Ind. Park, Hsin Chu Hsien, Taiwan,
R.O.C.

Product Registration Holder:

RIGEL PHARMA SDN. BHD.

3rd Floor, Apex Healthcare Building, No. 2,
Jalan SS 13/5, 47500 Subang Jaya, Selangor,
Malaysia.

Date of Revision: 13 February 2020

Version number	MY-BSN100-EI-2020.02.18
Revision date	2020.02.18
Dimensions (mm) Width * Height * Depth	140 x 200
PANTONE Matching System	