

FRONT

DISUF-B CREAM

VIDIS03-var2

DESCRIPTION

White, opaque cream

COMPOSITION

Fusidic Acid (Micronised) 2.0% w/w
Betamethasone valerate equivalent to Betamethasone 0.1% w/w

ACTIONS AND PHARMACOLOGY

Fusidic acid is an antibiotic with a bacteriostatic or bactericidal activity, mainly against Gram positive bacteria. Fusidic acid inhibits bacterial protein synthesis although, in contrast to drugs such as the macrolides or tetracyclines, it does not bind to the bacterial ribosomes, but inhibits a factor necessary for translocation of peptide subunits and elongation of the peptide chain. Betamethasone valerate is a topical steroid with rapid effect in inflammatory dermatoses. Even refractory conditions can often be treated successfully.

PHARMACOKINETICS

In vitro studies show that fusidic acid can penetrate intact human skin. The degree of penetration depends on factors such as the duration of exposure to fusidic acid and the condition of the skin. Betamethasone can be absorbed when given locally.

INDICATIONS

Disuf-B Cream is indicated in inflammatory dermatoses where bacterial infection is present or likely to occur. Inflammatory dermatoses include atopic eczema, discoid eczema, stasis eczema, seborrhoeic dermatitis, contact dermatitis, lichen simplex chronicus, psoriasis, discoid lupus erythematosus.

CONTRAINDICATIONS

- Topical corticosteroids are contraindicated in viral, tuberculous and fungal skin infections, perioral dermatitis, acne rosacea and ulcerative condition.
- Patients with hypersensitivity to any of the ingredients.

WARNING / PRECAUTIONS

- Long term continuous should be avoided.
- Caution should be exercised when the product is used near the eye, since topical steroid may cause glaucoma if the preparation gets into the eye.
- Steroid antibiotic combination should not be used for more than 7 days in the absence of any clinical improvement.

PREGNANCY AND LACTATION

Topical steroids should not be used extensively in pregnancy as fetal abnormalities have been observed in animals due to systemic absorption.

BACK

hovid

DISUF-B CREAM

MAIN SIDE/ADVERSE EFFECTS

- Hypersensitivity has rarely been reported.
- Prolonged and intensive treatment with topical steroids may cause local atrophic changes of the skin, such as striae, thinning and dilatation of the superficial blood vessels, particularly when occlusive dressing is used or when skin folds are involved.
- Systemic absorption, sufficient to produce hypercorticism/adrenal suppression, can occur with prolonged or extensive use, especially when used under occlusion/napkin.

OVERDOSE

Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects.

DOSAGE AND ADMINISTRATION

Apply sparingly to the affected area:

- Uncovered lesions : 2-3 times daily.
- Covered lesions : Less frequent application may be adequate.

Note: The information given here is limited. For further information consult your doctor or pharmacist.

Storage: Store below 25°C.

Presentation / Packing: Cream x 15 g aluminium tube.

Product Registration Holder / Manufactured by: HOVID Bhd.
121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar), 30010 Ipoh, Perak, Malaysia.

Revision date: November 2021.

hovid