

LIPROCT® SUPPOSITORY

Ingredient(s):

Each suppository contains:

Lidocaine	60mg
Hydrocortisone acetate	5mg
Zinc oxide	400mg

Pharmacology (Summary of Pharmacodynamic and Pharmacokinetic):

Lidocaine is a local anaesthetic of the amide type. It causes a reversible blockade of impulse propagation along nerve fibres by preventing the inward movement of sodium ions through the nerve membrane.

Hydrocortisone acetate belongs to the mild group of corticosteroids and is effective because of its anti-inflammatory and anti-pruritic action.

Zinc oxide is added as astringent agents.

Lidocaine is well-absorbed from the gastrointestinal tract, but little intact drug appears in the circulation because of biotransformation in the liver. Normally about 65% of the lidocaine is bound to plasma proteins. Lidocaine crosses the blood-brain and placental barriers, presumably by passive diffusion. The main elimination pathway of lidocaine is by liver metabolism. The primary route of lidocaine in human is dealkylation to form monoethylglycine-xylidine (MEGX) and glycinoxylidide (GX). Both of these metabolites may contribute to the therapeutic and toxic effects of lidocaine. Since GX has a longer half-life than lidocaine, accumulation may occur during long-term administration. Further metabolism occurs and metabolites are excreted in the urine with less than 10% of unchanged lidocaine. Renal impairment does not affect the clearance of lidocaine but accumulation of its active metabolites can occur and may lead to toxicity.

Less than 50% of hydrocortisone is absorbed following rectal application. When administered by topical application, particularly under an occlusive dressing or when the skin is broken, sufficient corticosteroid may be absorbed to give systemic effects. Corticosteroids in the circulation are extensively bound to plasma proteins, mainly to globulin and less to albumin. Only unbound hydrocortisone has pharmacological effects or is metabolized. Corticosteroids are metabolized mainly in the liver but also in the kidney, and are excreted in the urine.

Indication(s):

For the treatment of pain, itching and discomfort arising from irritated anorectal tissues e.g. haemorrhoids, pruritus ani, proctitis, milder forms of anal fissures, postoperative pain relief.

Dosage and Administration:

Insert one suppository morning and night and after each defecation. If defecation is painful, apply a few minutes before. The suppository is inserted just inside the anus.

A daily dose of 5 suppositories is well within safety limits. The duration of treatment may vary between ten days and three weeks. If the treatment is prolonged, a free interval can be recommended, especially if it is suspected that irritation due to lidocaine or hydrocortisone has occurred. If the local irritation disappears after the cessation of treatment, the possibility of sensitivity to lidocaine or hydrocortisone can be investigated, for example, by a patch test.

Contraindication(s):

Known hypersensitivity to local anaesthetics of the amide type or to other components of the product. Liproct should not be used in patients with untreated infections of bacterial, viral, pathogenic fungal or parasitic origin.

Side Effect(s) / Adverse Reaction(s):

Contact sensitivity to lidocaine has been reported after perianal use. Contact sensitivity may also occur after the use of topical hydrocortisone. In extremely rare cases amide-type local anaesthetic preparations have been associated with allergic reactions (in the most severe instances anaphylactic shock).

Precaution(s) / Warning(s):

1. Excessive dosage of lidocaine or short intervals between doses, may result in high plasma levels of lidocaine and serious adverse effects. Patients should be instructed to strictly adhere to recommended dosage.
2. The possibility of malignancy should be excluded before use.
3. If irritation or rectal bleeding develops, treatment should be discontinued and appropriate therapy instituted.
4. Systemic absorption of lidocaine may occur from the rectum, and large doses may result in CNS side-effects. On rare occasions convulsions have occurred in children.
5. Prolonged and excessive use of hydrocortisone may produce systemic corticosteroid effects or local effects such as skin atrophy. With the recommended dosage systemic effects of hydrocortisone are unlikely.

Interaction with Other Medicaments:

Lidocaine should be used with caution in patients receiving antiarrhythmic drugs, since the toxic effects are additive.

With large doses of lidocaine, consideration should be given to the risk of additional systemic toxicity in patients receiving other local anaesthetics or agents structurally related to local anaesthetics.

Pregnancy and Lactation:

It is reasonable to assume that a large number of pregnant women and women of child-bearing age have been given lidocaine and hydrocortisone. No specific disturbances to the reproductive process have so far been reported, e.g. no increased incidence of malformations.

Lidocaine and hydrocortisone acetate are excreted into breast milk but in such small quantities that adverse effects on the child are unlikely at therapeutic doses.

Symptoms and Treatment for Overdosage, and Antidote(s):

Lidocaine can cause acute toxic effects if high systemic levels occur due to rapid absorption or overdosage. With the recommended doses, toxic effects have not been reported. However, should systemic toxicity occur, the signs are anticipated to be similar in nature to those following the administration of local anaesthetics by other routes.

Local anaesthetic toxicity is manifested by symptoms of nervous system excitation and, in severe cases, central nervous and cardiovascular depression.

Severe neurological symptoms (convulsions, CNS depression) must be treated symptomatically by respiratory support and the administration of anticonvulsive drugs.

Shelf-Life:

2 years from the date of manufacture.

Storage Condition(s):

Store at temperature below 25°C. Protect from light and moisture.

Product Description and Packing(s):

White to off-white spindle suppositories.

Case of 10's x 1, 10's x 2 and 10's x 10.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679