

**USER INFORMATION
READ CAREFULLY**

RinsCap® CG
(CHLORHEXIDINE GLUCONATE 0.05% w/v IRRIGATION SOLUTION)
Sterile Antiseptic Solution

Product Description:

Clear, Blue, Sterile Irrigation Solution.

Composition:

The Solution Contains: Chlorhexidine Gluconate 0.05% w/v

Pharmacodynamics:

Chlorhexidine is a disinfectant which is effective against a wide range of vegetative Gram-positive and Gram-negative bacteria: it is more effective against Gram-positive than Gram-negative bacteria, some species of *Pseudomonas* and *Proteus* being less susceptible. The wide range of organisms against which chlorhexidine is active explains the rationale for presenting it in a solution for swabbing wounds and burns and in obstetrics.

Pharmacokinetics:

Chlorhexidine is poorly absorbed from the gastrointestinal tract and skin.

Indication:

Chlorhexidine Gluconate is a potent antibacterial agent for general antiseptic purposes. It is bactericidal to a broad spectrum of organisms. RinsCap® CG is recommended for use in obstetrics and for swabbing burns and wounds.

Recommended Dosage:

Rinse the target area with clean water. Gently wash the area with the minimum amount of RinsCap® CG. The solution is used undiluted. After rinsing with undiluted RinsCap® CG, rinse the target area again with clean water.

Route of Administration:

For external use

Contraindications:

RinsCap® CG should not come into contact with the brain, eyes, meninges or middle ear. The use of RinsCap® CG is also contraindicated in patients who have shown hypersensitivity to chlorhexidine.

Warnings and Precautions:

For external use only. The solution is hypotonic and haemolytic. Use undiluted. Do not mix with detergents or other chemicals. Do not heat this unit in excess of 80°C. Do not use unless seal is intact.

RinsCap® CG contains chlorhexidine. Chlorhexidine is known to induce hypersensitivity, including generalised allergic reactions and anaphylactic shock. The prevalence of chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare. RinsCap® CG should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine.

If any signs of symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate develop, immediately stop using the product. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Interactions With Other Medicaments:

Hypochlorite bleaches may cause brown stains to develop in fabrics, which have previously been in contact with RinsCap® CG.

Statement On Usage During Pregnancy and Lactation:

Although there are no adverse reports for this product in pregnant and lactating mothers, as with all medicines, care should be exercised when administering the product to pregnant or lactating women.

Adverse Effects/Undesirable Effects:

Idiosyncratic skin reactions and generalized allergenic reactions to chlorhexidine can occur, but these are rare.

Immune system disorders

Frequency not known: Hypersensitivity including anaphylactic shock.

Overdose and Treatment:

The main consequence of ingestion is mucosal irritation and systemic toxicity is rare due to minimal absorption from the gastrointestinal tract. Haemolysis has been reported after accidental intravenous administration. Gastric lavage with demulcents has been suggested for acute ingestion.

Storage Condition:

Do not store above 30°C. Protect from light.

Shelf Life:

3 years from manufacturing date in the proposed storage condition.
Do not use after expiry.

Dosage Form and Packaging Available:

500 mL x 20 PP bottle per box
1000 mL x 10 PP bottle per box

Manufacturer/Product Registration Holder:

 **AIN MEDICARE SDN. BHD.**
Lot 4933, 4934 & PT2464, Jalan 6/44,
Kaw. Perindustrian Pengkalan Chepa 2,
16100 Kota Bharu, Kelantan, MALAYSIA

Date of Revision: 01.08.2020