



Pregnancy and lactation

Usage in Pregnancy: There are no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always, predictive of human response, this drug should be used during pregnancy only if the potential benefit to the patient justifies the potential risk to the fetus.

Usage in Nursing Mothers: Gabapentin is excreted in human milk. Because the effect on the nursing infant is unknown, caution should be exercised when gabapentin is administered to nursing mother.

Gabapentin should be used in nursing mothers only if the benefits clearly outweigh the risks.

Side effects

The adverse reactions in epilepsy (adjunctive and monotherapy) and neuropathic pain have been provided in a single list below by class and frequency (very common > 1/10), common >1/100, <1/10),

uncommon >1/1000, <1/100) and rare >1/10,000; <1/1,000).

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Infections and infestations

Very Common: Viral infection

Common: Pneumonia, respiratory infection, urinary

tract infection, infection, otitis media

Blood and the lymphatic system disorders

Common: leucopenia

Rare: thrombocytopenia

Immune system disorders

Rare: allergic reactions (e.g. urticaria)

Metabolism and Nutrition Disorders

Common: anorexia, increased appetite

Psychiatric disorders

Common: hostility, confusion and emotional lability, depression, anxiety, nervousness, thinking abnormal

Rare: hallucinations

Nervous system disorders

Very Common: somnolence, dizziness, ataxia,

Common: convulsions, hyperkinesias, dysarthria, amnesia, tremor, insomnia, headache, sensations such as paresthesia, hypaesthesia, coordination abnormal, nystagmus, increased, decreased, or absent reflexes

Rare: movement disorders (e.g. choreoathetosis, dyskinesia, dystonia)

Eye disorders

Common: visual disturbances such as amblyopia, diplopia

Ear and Labyrinth disorders

Common: vertigo

Rare: tinnitus

Cardiac disorders

Rare: palpitations

Vascular disorder

Common: hypertension, vasodilatation

Respiratory, thoracic and mediastinal disorders

Rare: Respiratory depression

Post-marketing experience: Dysphagia

Common: dyspnoea, bronchitis, pharyngitis, cough, rhinitis

Gastrointestinal disorders

Common: vomiting, nausea, dental abnormalities, gingivitis, diarrhea, abdominal pain, dyspepsia, constipation, dry mouth or throat, flatulence

Rare: pancreatitis

Hepatobiliary disorders

Rare: hepatitis, jaundice

Skin and subcutaneous tissue disorders

Common: facial oedema, purpura most often described as bruises resulting from physical trauma, rash, pruritus, and acne

Rare: Stevens-Johnson syndrome, angioedema, erythema multiforme, alopecia

Musculoskeletal, connective tissue and bone disorders

Common: arthralgia, myalgia, back pain, twitching

Renal and urinary disorders

Common: incontinence

Rare: acute renal failure

Reproductive system and breast disorders

Common: impotence

Symptoms and Treatment of over dose

Acute, life-threatening toxicity may not occur with Gabapentin overdoses of up to 49 grams. Symptoms of the overdoses included dizziness, double vision, slurred speech, drowsiness, lethargy and mild diarrhoea. All patients recovered fully with supportive care. Reduced absorption of Gabapentin at higher doses may limit drug absorption at the time of overdosing and, hence, minimise toxicity from overdoses.

Although gabapentin can be removed by hemodialysis, based on prior experiment, hemodialysis may be indicated.

An oral lethal dose of gabapentin was not identified in mice and rats given doses as high as 8000 mg/kg. Signs of acute toxicity in animals included ataxia, labored breathing, ptosis, hypoactivity, or excitation.

Storage condition

Do not store above 30°C.

Shelf life

24 months.

Presentation:

ONEGABA 300 MG CAPSULES:

3 blisters of 10 capsules each are packed in a printed carton along with a package insert.

10 blisters of 10 capsules each are packed in a printed carton along with a package insert.

ONEGABA 400 MG CAPSULES:

3 blisters of 10 capsules each are packed in a printed carton along with a package insert.

The effects of long-term (greater than 36 weeks) gabapentin therapy on learning, intelligence, and development in children and adolescents has not been adequately studied. The benefits of prolonged therapy must therefore be weighed against the potential risks of such therapy.

Manufactured By:



Aurobindo Pharma Limited (Unit III),

Survey No. 313 & 314, Bachupally,

Bachupally Mandal, Medchal-Malkajigiri District,

Telangana State, INDIA.

DATE OF PREPARATION OF THIS LEAFLET

April 2025