

CP Cough Syrup

VICPCNa-PC (MY)

DESCRIPTION

Brownish liquid with vanilla flavour

COMPOSITION

Each 5 ml contains:

Carbocisteine	100.0 mg
Promethazine HC/	2.5 mg

ACTIONS AND PHARMACOLOGY

Promethazine, a phenothiazine derivative, is a sedating antihistamine with antimuscarinic, significant sedative, and some serotonin-antagonist properties. It is usually given as the hydrochloride or teoclate. Promethazine Hydrochloride has anticholinergic actions of most antihistamines which provide a drying effect on the nasal and oral mucosa. It is commonly use in the preparation for symptomatic treatment of coughs and the common cold.

Carbocisteine is used for its mucolytic activity in respiratory disorders associated with productive cough. It is usually given by mouth as the sodium or lysine salts.

PHARMACOKINETICS

Promethazine is well absorbed after oral administration. Peak plasma concentrations have been observed 2 to 3 hours after administration. Onset of action occurs 15 to 60 minutes after oral administration and may have the duration of action from 4 to 12 hours. Promethazine crosses the blood-brain barrier and the placenta, and is distributed into breast milk. Values ranging from 76 to 93% have been reported for plasma-protein binding. It is excreted slowly via the urine and bile. Elimination half-lives of 5 to 14 hours have been reported.

Carbocisteine is rapidly and well absorbed from gastrointestinal tract with peak plasma concentrations occurring 90 to 120 minutes after oral dose. It appears to penetrate into lung tissue and respiratory mucus. Carbocisteine is excreted in the urine as unchanged drug and metabolites. Acetylation, decarboxylation, and sulfoxidation have been identified as the major metabolic pathways. Sulfoxidation may be governed by genetic polymorphism.

INDICATIONS

- Symptomatic treatment of troublesome, nonproductive cough, particularly occurring at night.
- This product contains Promethazine HC/. It should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression.

CONTRAINDICATIONS

- CP Cough Syrup is contraindicated in patients with gastroduodenal ulcer.
- This medication should not be used in patients with sensitivity to Promethazine HC/ and Carbocisteine.
- The presence of other medical problems may affect the use of this medication. A doctor should be consulted before using this medicine if any other medical problems exist, especially: bladder neck obstruction, symptomatic prostatic hypertrophy, urinary retention, bone marrow depression, cardiovascular disease, coma, glaucoma, jaundice, Reye's syndrome, asthma, chronic bronchitis, productive cough, emphysema, diabetes, hepatic function impairment and respiration depression.
- Use of this medicine is inadvisable in combination with sultopride.
- Contraindicated in children below 2 years of age.

PRECAUTIONS

Caution should be exercised to patients sensitive to promethazine or to other phenothiazine medications:

- **For pediatrics:** Use of promethazine is not recommended in newborn or premature infants because this age group has an increased susceptibility to anticholinergic side effects, such as central nervous system (CNS) excitation, and an increased tendency toward convulsions.
- **For adolescents:** Promethazine should not be used in adolescents with signs and symptoms suggestive of Reye's syndrome since extrapyramidal symptoms may occur.
- **For geriatrics:** Dizziness, sedative, confusion, hypotension and paradoxical reaction may be more likely to occur in geriatric patients taking promethazine.
- Prolonged use of promethazine may decrease or inhibit salivary flow. Involuntary orofacial muscle movements may result from extrapyramidal effects.

Carbocisteine which may disrupt the gastric mucosal barrier should be used in caution in patients with a history of peptic ulceration.

WARNING

This medicine may cause drowsiness and dizziness, less alert than normal or to feel a false sense of well being. Patients should know how they react to this medicine before they drive, use machines or doing anything that could be dangerous if they are dizzy or are not alert and clearheaded. Children should be supervised to avoid potential harm in bike riding or other dangerous activities. When used for treatment of cough and cold: It should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression. To be used with caution and doctor's/pharmacist's advice in children 2 to 6 years of age.

PREGNANCY AND LACTATION

Safety of use has not been established. Risk benefits should be considered when using on pregnant or lactating mothers.

DRUG INTERACTIONS

Concurrent use of CP Cough Syrup and the following drugs must be avoided or if necessary, use with care:

- Alcohol
- Other central nervous system depressants (sedatives antidepressants, barbiturates, clonidine and related substances, hypnotics, morphine derivatives (analgesics and antitussives), methadone, neuroleptics, anxiolytics)
- Atropine and other atropine substances (imipramine antidepressants, anticholinergic antiparkinsonians, atropine antispasmodics, disopyramide, phenothiazine neuroleptics)
- Sultopride

MAIN SIDE/ADVERSE EFFECTS

The side effects with this medication appear to be rare and may include drowsiness, blood dyscrasias, nausea, dizziness, gastrointestinal bleeding, gastric discomfort, and other common side effects of sedating antihistamines like headache, confusion, difficult or painful urination, dryness of mouth, nose or throat, hypotension, increased sweating, loss of appetite, paradoxical reaction, photosensitivity, ringing or buzzing in ears, skin rash and tachycardia.

OVERDOSE

- **Symptoms of overdose:** Convulsions, consciousness disorders, coma.
- **Treatment for overdose:** Treatment is symptomatic and should be initiated in a specialized unit.

DOSAGE AND ADMINISTRATION

For oral use.

Adult	: 3 to 4 tablespoons daily
Children 12-15 years	: 6 to 9 teaspoons daily
Children 30 months-12 years	: 4 to 6 teaspoons daily
Children 24-30 months	: 3 to 4 teaspoons daily

This product contains Promethazine HCl. It should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression.

This medication should be taken with food. Patients are advised to use a calibrated measuring device as 5 ml is a rough approximation of one teaspoon and different teaspoons differ in their volume.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage:
Store below 30°C. Protect from light.

Presentation/Packing:
120 ml per amber glass bottle

Product Registration Holder/
Manufactured by: HOVID Bhd.
121, Jalan Tunku Abdul Rahman
(Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Revision date: March 2018