

CETIHIS Syrup**ANTI-HISTAMINE (H₁-RECEPTOR)**

Each 5 mL contains:
Cetirizine dihydrochloride 5 mg

✧ **PRODUCT DESCRIPTION:**

Clear, colorless syrup with banana-grape flavor

✧ **MECHANISM OF ACTION:****Pharmacology**

In animals, CETIHIS has been shown to be an anti-H₁ agent devoid of any significant anticholinergic or antiserotonin effects. At active doses, it does not induce sedation or behavioral changes.

CETIHIS will inhibit certain effects produced by exogenous histamine. CETIHIS also inhibits the effects produced by endogenous histamine released *in vivo*. It inhibits the cutaneous reaction induced by Vasoactive Intestinal Polypeptide and substance P, neuropeptides.

CETIHIS inhibits the histamine-induced "early" phase of the allergic reaction. It also significantly reduces the migration of inflammatory cells e.g., eosinophils and the release of mediators associated with the "late" allergic response.

CETIHIS reduces bronchial hyper reactivity to histamine in the asthmatic patient. It also reduces the allergic reaction induced by specific allergens. These effects are obtained without any central effects.

Pharmacokinetics

Peak blood levels reached between 30 and 60 minutes after administration of 10 mg CETIHIS. It is strongly bound to plasma proteins.

✧ **INDICATION:**

Adults and children of 2 years and above: symptomatic treatment of seasonal allergic rhinitis, perennial allergic rhinitis and urticaria of allergic origin.

✧ **DOSAGE AND ADMINISTRATION:**

The solution can be swallowed as such.

Route of Administration

For oral use.

Adults

10 mg (10 mL of oral solution) once daily.

A 5 mg starting dose (5 mL of oral solution) may be proposed if this leads to satisfactory control of the symptoms.

Children

Children aged from 2 to 6 years

2.5 mg (2.5 mL of oral solution bid) twice daily.

Children aged from 6 to 12 years

5 mg (5 mL of oral solution) twice daily.

Children over 12 years of age

10 mg (10 mL of oral solution) once daily.

Elderly

Data does not suggest that the dose needs to be reduced in elderly subjects provided that the renal function is normal.

Patients with moderate to severe renal impairment

Since Cetirizine is mainly excreted via renal route, in cases no alternative treatment can be used, the dosing intervals must be individualized according to renal function.

Refer to the following table and adjust the dose as indicated.

$$CL_{Cr} = \frac{[140 - \text{age (years)}] \times \text{weight (kg)}}{72 \times \text{serum creatinine (mg/dL)}} \times 0.85 \text{ for women}$$

Dosing adjustments for adult patients with impaired renal function:

Group	Estimated Glomerular Filtration Rate (eGFR) (mL/min)	Dosage and frequency
Normal renal function	≥ 90	10 mg once daily
Mild decrease renal function	60 – < 90	10 mg once daily
Moderately decreased renal function	30 – < 60	5 mg once daily
Severely decreased renal function	15 – < 30 not requiring dialysis treatment	5 mg once every 2 days
End-stage renal disease	< 15 requiring dialysis treatment	Contraindicated

In pediatric patients suffering from renal impairment, the dose will have to be adjusted on an individual basis taking into account the renal clearance, age and body weight of the patient.

Patients with hepatic impairment

No dose adjustment is needed in patients with solely hepatic impairment.

Patients with hepatic impairment and renal impairment

Dose adjustment is recommended (see *Patients with renal impairment above*).

✧ **WARNING AND PRECAUTION:****Alcohol**

At therapeutic doses, no clinically significant interactions have been demonstrated with alcohol (for a blood alcohol level of 0.5 g/L). Nevertheless, precaution is recommended if alcohol is taken concomitantly (see *Section Drug Interaction*).

Increased risk of urinary retention

Caution should be taken in patients with predisposition factors of urinary retention (e.g. spinal cord lesion, prostatic hyperplasia) as Cetirizine may increase the risk of urinary retention (see *Section Side Effect*).

Patients at risk of convulsions

Caution in epileptic patients and patients at risk of convulsions is recommended.

Skin reactions

Pruritus and/or urticaria may occur when Cetirizine is stopped, even if those symptoms were not present before treatment initiation (see *Section Side Effect*). In some cases, the symptoms may be intense and may require treatment to be restarted. The symptoms should resolve when the treatment is restarted.

Allergy skin tests

Allergy skin tests are inhibited by antihistamines and a wash-out period of 3 days is recommended before performing them.

Food

The extent of absorption of Cetirizine is not reduced with food, although the rate of absorption is decreased.

Activities requiring mental alertness

In clinical trials, the occurrence of somnolence has been reported in some patients taking Cetirizine. Due caution should therefore be exercised when driving a car or operating potentially dangerous machinery.

Pregnancy and Lactation**Fertility**

Limited data is available on human fertility but no safety concern has been identified.

Animal data show no safety concern for human reproduction.

Product	Code No.	Dimension	Packaging Type	Thickness
CETIHIS Syrup	ICMY 0053	W 17 x L 25 cm.	Wood Free Paper (กระดาษปลอดไม้)	60 g (0.08 mm.)
 Designed by: (PDM, ASPD, PDC-M, PDC-A, PDC-H, PDC-A) Checked by: (PDM/ASPD) Approved by: (MD (Only Pattern and Colors))				
Approved by: PLC: LRA: IRA: QCC-PM: CUSTOMER/SALES DEPT.:				

Pregnancy
Caution should be exercised when prescribing to pregnant women. For Cetirizine prospectively collected data on pregnancy outcomes do not suggest potential for maternal or fetal/embryonic toxicity above background rates.
Lactation
Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/fetal development, parturition or postnatal development.
Cetirizine passes into breast milk. A risk of side effects in breastfed infants cannot be excluded. Caution should be exercised when prescribing Cetirizine to lactating women. Cetirizine is excreted in human milk at concentrations representing 25% to 90% of those measured in plasma, depending on sampling time after administration.

❖ **SIDE EFFECT:**
Post Marketing Data
Adverse drug reactions (ADRs) are listed below by MedDRA system organ class and by frequency.
Frequencies are defined as: Very common (≥1/10); Common (≥1/100 to <1/10); Uncommon (≥1/1,000 to <1/100); Rare (≥1/10,000 to <1/1,000); Very rare (<1/10,000); Not known (cannot be estimated from the available data).

System Organ Class	Frequency	Undesirable Effect
Blood and lymphatic system disorders	Very rare	Thrombocytopenia
Immune system disorders	Rare	Hypersensitivity
	Very rare	Anaphylactic shock
Metabolism and nutrition disorders	Not known	Increased appetite
Psychiatric disorders	Uncommon	Agitation
	Rare	Aggression, confusion, depression, hallucination, insomnia
	Very rare	Tics
	Not known	Suicidal ideation, nightmare
Nervous system disorders	Uncommon	Paresthesia
	Rare	Convulsions
	Very rare	Dysgeusia, dyskinesia, dystonia, syncope, tremor
	Not known	Amnesia, memory impairment
Eye disorders	Very rare	Accommodation disorder, blurred vision, oculogyric crisis
Ear and labyrinth disorders	Not known	Vertigo
Cardiac disorders	Rare	Tachycardia
Gastrointestinal disorders	Uncommon	Diarrhea
Hepatobiliary disorders	Rare	Hepatic function abnormal (transaminases increased, blood bilirubin increased, blood alkaline phosphatase increased, gamma-glutamyl transferase increased)
	Not known	Hepatitis
Skin and subcutaneous tissue disorders	Uncommon	Pruritus, rash
	Rare	Urticaria
	Very rare	Angioedema, fixed drug eruption
	Not known	Acute generalized exanthematous pustulosis (AGEP)
Musculoskeletal and connective tissue disorders	Not known	Arthralgia, myalgia
Renal and urinary disorders	Very rare	Dysuria, enuresis
	Not known	Urinary retention (see Section Warning and Precaution)
General disorders and administration site conditions	Uncommon	Asthenia, malaise
	Rare	Oedema
Investigations	Rare	Weight increased

Skin reactions occurring after discontinuation of Cetirizine
After discontinuation of Cetirizine, pruritus (intense itching) and/or urticaria have been reported (see Section Warning and Precaution).

❖ **CONTRAINDICATION:**
Cetirizine is contraindicated in:
• hypersensitivity to any of the constituents of this formulation, to Hydroxyzine or to any Piperazine derivatives
• patients with end-stage renal disease with eGFR (Estimated Glomerular Filtration Rate) below 15 mL/min.
❖ **DRUG INTERACTION:**
Due to the pharmacokinetic, pharmacodynamic and tolerance profile of Cetirizine, no interactions are expected with this antihistamine. Neither pharmacodynamic nor significant pharmacokinetic interaction was reported in drug-drug interactions studies performed, notably with Pseudoephedrine or Theophylline (400 mg/day).
Alcohol and other CNS depressants
In sensitive patients, the concurrent use of alcohol or other CNS depressants may cause additional reductions in alertness and impairment of performance, although Cetirizine does not potentiate the effect of alcohol (0.5 g/L blood levels) (see Section Warning and Precaution).

❖ **OVERDOSE AND TREATMENT:**
Symptoms and signs
Symptoms observed after an important overdose of Cetirizine are mainly associated with CNS effects or with effects that could suggest an anticholinergic effect. Adverse events reported after an intake of at least 5 times the recommended daily dose are: confusion, diarrhea, dizziness, fatigue, headache, malaise, mydriasis, pruritus, restlessness, sedation, somnolence, stupor, tachycardia, tremor, and urinary retention.
Treatment
There is no known specific antidote to Cetirizine.
Should overdose occur, symptomatic or supportive treatment is recommended.
Cetirizine is not effectively removed by hemodialysis.
Management should be as clinically indicated or as recommended by the national poisons centre, where available.

❖ **STORAGE:**
Store at temperature of not more than 30°C.

❖ **DOSAGE FORM AND PACKAGING AVAILABLE:**
Bottle 5 mg/5 mL, 60 mL Syrup

❖ **DATE OF REVISION:**
July 18, 2025

Manufactured by:
UNISON LABORATORIES CO., LTD.
39 Moo 4, Klong Udomcholjorn, Muang Chachoengsao,
Chachoengsao 24000 Thailand

C-MY/180725-16 (AR)
(MIMS ANNUAL Malaysia 2005; ^ ZYRTEC_Feb. 2024)

ICMY 0053

PM Specification	Product	Code No.	Dimension	Packaging Type	Thickness
	CETIHIS Syrup	ICMY 0053	W 17 x L 25 cm.	Wood Free Paper (กระดาษปลอดไม้)	60 g (0.08 mm.)
Designed by: (PDM, ASD, PDC-M, PDC-A, PDO-M, PDC-A) Checked by: (PDM/ASPD) Approved by: (IMD (Only Pattern and Colors))					
Approved by: (Date)	PLC: (Date)	LRA: (Date)	IRA: (Date)	QCC-PM: (Date)	CUSTOMER/SALES DEPT: (Date)