


ZEFXON IV INJECTION 40 mg



COMPOSITION

Each pack contains one vial and one ampoule :
Each vial contains : Omeprazole sodium 42.6 mg equivalent to Omeprazole 40 mg
Each ampoule contains : 10 ml of solvent for reconstitution

PRODUCT DESCRIPTION

Drug Substance

Dry, homogeneous, white crystalline powder

Solvent for reconstitution

Clear, colorless solution

Product after reconstitution

Clear and free from visible evidence of particles.

PHARMACODYNAMICS

Zefxon injection (Omeprazole) is the proton/ acid pump inhibitor which reduces gastric acid secretion by specific action ; inhibition of the enzyme H⁺, K⁺ - ATPase (The proton pump) in the parietal cell, the final step of acid secretion ; thus inhibition acid secretion occurs.

PHARMACOKINETICS

Distribution

The apparent volume of distribution in healthy subjects is approximately 0.3 L/kg and a similar value is also seen in patients with renal insufficiency. In the elderly and in patients with hepatic insufficiency, the volume of distribution is slightly decreased. The plasma protein binding of omeprazole is about 95%.

Metabolism and excretion

The average half-life of the terminal phase of the plasma concentration-time curve following IV administration of omeprazole is approximately 40 minutes ; the total plasma clearance is 0.3 to 0.6 L/min. There is no change in half-life during treatment.

Omeprazole is completely metabolised by the cytochrome P450 system, mainly in the liver.

No metabolite has been found to have any effect on gastric acid secretion. Almost 80% of an intravenously given dose is excreted as metabolites in the urine, and the remainder is found in the faeces, primarily originating from biliary secretion.

Elimination of omeprazole is unchanged in patients with reduced renal function. The elimination half-life is increased in patients with impaired liver function, but omeprazole has not shown any accumulation with once daily oral dosing.

INDICATIONS

Zefxon injection is indicated for alternative use to oral medication in

- Duodenal ulcer
- Gastric ulcer
- Reflux oesophagitis
- Zollinger-Ellison syndrome

DOSAGE AND ADMINISTRATION

Zefxon injection administration once daily is recommended where oral medication is inappropriate. This produces a mean decrease over 24 hours of approximately 90%. In patients with Zollinger-Ellison syndrome the recommended initial dosage of Zefxon injection is 60 mg once daily. Higher daily doses may be required and the dose should be divided and given twice daily.

Direction for Reconstitution

Dissolve Zefxon injection 1 vial with the solvent provided to get omeprazole concentration of 4 mg/mL.

The IV solution should be used within 4 hours after reconstitution. Do not use if any particles are present in the reconstituted solution.

Incompatibilities

None known when instructions in Dosage and Administration are followed.

Use in children :

Safety and efficacy of omeprazole in children have not been established.

Use in elderly :

Normal adult dosage is recommended.

Dosage in Renal Impairment :

Although pharmacokinetics may be altered in patients with renal impairment, dosage adjustment dose not appear necessarily in patients with such impairment.

Dosage in Hepatic Impairment :

Dosage adjustment should be considered in patients with hepatic impairment. Some clinicians recommend that such patients with hepatic dysfunction receiving dosage exceeding 20 mg daily should be monitored closely for possible adverse effects.

Pregnancy and lactation :