



VITAMIN C TABLET

Each tablet contains:
Ascorbic Acid 100mg

Indications:

As a dietary supplement.

Contraindications:

It is contraindicated in patients who are sensitive to Ascorbic Acid or the other ingredients in the product.

Side effects/adverse reactions:

Ascorbic Acid is usually well tolerated. However, allergic responses to it presenting as eczema, urticaria or asthma have been reported.

Large doses may cause diarrhoea and the formation of renal calcium oxalate calculi.

Precautions / Warnings:

Ascorbic Acid could be partly metabolised to oxalic acid. Average reported production rates were 1mg oxalate from 1g, 12mg from 4g, and 69mg from 9g, but with wide individual variations. Some patients would have a high risk of renal calcification during treatment with high doses of ascorbic acid.

Tolerance may be induced in patients taking high doses. When daily ascorbic acid intake significantly exceeded 250mg for prolonged periods, the rate of clearance from the body increased so that blood concentrations did not increase with large doses. Dependence on larger than normal maintenance doses could occur and deficiency might occur if treatment was withdrawn.

Ascorbic Acid can cause chemical interference in laboratory tests for blood or urine creatinine, glucose and uric acid.

It is reported that mean plasma ethinyl-oestradiol concentrations were increased by 16.3 and 47.6% six and 24 hours respectively after ingestion of Ascorbic Acid. The overall effect of a large Vitamin C supplement in patients on contraceptives, is the conversion of a low oestrogen oral contraceptive into a high-dose contraceptive.

Ascorbic Acid taken concomitantly with fluphenazine may necessitate an increase in dosage of fluphenazine.

Ascorbic Acid in large doses might obscure the diagnosis of gout or in some patients it might precipitate acute gouty arthritis.

Dosage:

Oral administration.

Adults : 1-2 tablets 2 to 4 times daily.

Children : 1-3 tablets daily.

Drug interactions:

Ascorbic Acid has to be given with extreme caution to patients with excess tissue iron; when used with desferrioxamine the need for Ascorbic Acid should be clearly established by measuring iron excretion before and after supplementation and the oral dose of Ascorbic Acid should be administered 1-2 hour after infusion has been started.

The mean plasma-ethinyl-oestradiol concentration is increased with concomitant administration of a large supplement of Vitamin C and an oral contraceptive containing ethinyl-oestradiol.

Increase in manic behaviour necessitating increase in dosage with fluphenazine followed treatment with Ascorbic Acid to reduce Vitamin C deficiency.

Symptoms and Treatment for overdosage and antidote(s):

Adverse reactions arising from overdosage of Ascorbic Acid appears remote. Vitamin C is water soluble, it is not significantly stored in the body. It is very well tolerated. Very high doses may cause gastro-intestinal discomfort and diarrhoea.

Subjects have been reported to tolerate doses up to 9g in dosage regimens of once and thrice daily without any adverse effects.

Pack sizes:

A bottle of 80, 100, 300 and

(For Export only)1000 tablets.

Alu-alu strip pack: 10 x 10, 50 x 10 and 100 x 10 tablets per strip.

Storage conditions: Store at or below 30°C. Protect from light.

Shelf-life: 3 years.

Description: Round, orange, sugar-coated tablet with no markings or embossment.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Manufacturer:

Sunward Pharmaceutical Sdn. Bhd.

No. 9, 11 & 17, Jalan Kempas 4,

Taman Perindustrian Tampoi Indah,

81200 Johor Bahru, Johor, Malaysia.

TL707B-R6
Revised Date: 26/03/2018