

For use only a Registered Medical Practitioner
or Hospital or a Laboratory only.

Package Insert

Quace

ECZEMA CREAM

Clobetasol Propionate Cream USP, 0.05%

Composition

Clobetasol Propionate USP 0.05% w/w
Chlorocresol BP (as preservative) 0.10% w/w
in a cream base q.s.

Description

White, smooth odourless cream free from any
grittiness and foreign matter.

Pharmacodynamics

Clobetasol propionate is a highly active corticosteroid
with topical anti-inflammatory activity. The major
effect of clobetasol propionate on skin is a non-
specific anti-inflammatory response, as a result of
vasoconstriction and decrease in collagen synthesis.

Pharmacokinetics

Clobetasol is absorbed through the skin, particularly
in denuded areas. Its penetration through the skin can
be increased by means of an occlusive dressing. The
absorbed clobetasol is metabolished in the liver and
excreted in the urine.

Indication

Clobetasol Propionate is a potent topical
corticosteroid indicated for short courses treatment of
more resistant dermatoses such as psoriasis
(excluding widespread plaque psoriasis), recalcitrant
eczemas, lichen planus, discoid lupus erythematosus,
and other conditions which do not respond
satisfactorily to less active steroids

Recommended Dose

Apply sparingly to the affected area once or twice daily
until improvement occurs; normally within a few days
in the more responsive conditions. Discontinue the
therapy when control is achieved. If a longer course is
required, it is recommended that treatment should not
be continued for more than 4 weeks without the

patient's condition being reviewed. Repeated short
courses of the Clobetasol Propionate may be used to
control exacerbations. A less potent preparation
should be used if continuous steroid treatment is
necessary. In vary resistant lesions, especially where
there is hyperkeratosis, the anti-inflammatory effect of
Clobetasol Propionate can be enhanced by occluding
the treatment area with polythene film. Overnight
occlusion is usually adequate to bring about
satisfactory response. Thereafter improvement can
usually be maintained by application without
occlusion

Mode Of Administration

For external use only.

Contraindication

Hypersensitivity to clobetasol propionate, to other
corticosteroids, or to any of the components of Quace
eczema cream is a contraindication to the use of this
product.

Warning and Precautions

Systemic absorption of topical corticosteroids should
always be considered. It should not be applied with an
occlusive dressing to large areas of the body. Long-
term continuous topical therapy should be avoided,
especially in children. If Clobetasol Propionate is
required for use in children, it is recommended that the
treatment should be reviewed weekly, so that the least
potent corticosteroid that will control the disease can
be selected. It should be noted that, the infant's napkin
might act as an occlusive dressing. The face, more
than other areas of the body, may exhibit atrophic
changes after prolonged treatment with potent topical
corticosteroids. If applied to the eyelids, care is
needed to ensure that the preparation does not enter
the eyes as might cause corneal ulcers, raised intra-
ocular pressure and reduced visual function. Any
spread of infection requires withdrawal of topical
corticosteroid therapy and systemic administration of
appropriate antimicrobial product. Topical
corticosteroids may be hazardous in psoriasis for a
number of reasons including rebound relapses,
development of tolerance, risk of generalized pustular
psoriasis and development of local and systemic
toxicity due to impair barrier function of the skin. If
used in psoriasis careful patient supervision is
important.

Drug Interactions

Potentially hazardous interactions
No hazardous interactions have been reported
Potentially useful interactions
No such interactions are reported, although
clobetasol topical applications can be combined with
an antibiotic

Pregnancy and Lactation

There is inadequate evidence of safety in human
pregnancy. Topical administration of corticosteroids
to pregnant animal can cause abnormalities of foetal
development. The relevance of this finding to human
beings has not been established, however, topical
steroids should not be used extensively in pregnancy
such as in large amounts for prolonged periods.
The safe use of topical corticosteroids during lactation
has not been established.

Side Effects

QUACE ECZEMA Cream is generally well tolerated
when used for two-week treatment periods.
The most frequent adverse reactions reported to
clobetasol propionate cream have been local and
have included burning sensation and stinging
sensation in approximately 1% of the patients. Less
frequent adverse reactions were itching, skin atrophy,
and cracking and fissuring of the skin.
The following local adverse reactions are reported
infrequently when topical corticosteroids are used as
recommended. These reactions are listed in an
approximately decreasing order of occurrence:
burning, itching, irritation, dryness, folliculitis,
hypertrichosis, acne form eruptions,
hypopigmentation, perioral dermatitis, allergic
contact dermatitis, maceration of the skin, secondary
infection, skin atrophy, striae, and miliaria. Systemic
absorption of topical corticosteroids has produced
reversible. HPA axis suppression, manifestations of
Cushing's syndrome, hyperglycemia and glucosuria
in some patients. In rare instances, treatment (or
withdrawal of treatment) of psoriasis with
corticosteroids is thought to have exacerbated the
disease or provoked is postural form of the disease, so
careful patient supervision is recommended

Symptoms And Treatment of Overdose

Acute overdosage is unlikely to occur, however, in
case of chronic overdosage or misuse the features of
hypercorticism may appear and topical
corticosteroids should be discontinued.

Storage Condition

Store at a temperature not exceeding 30°C.
Do not freeze.

Shelf Life

24 months from the date of manufacture.

Presentation:

15g aluminum collapsible tube and flowerpot type
white HDPE cap packed in a unit printed duplex board
carton.

Date of revision : Jan. 2021

Mfg.Lic No. : 28/9/96
Reg. No. : MAL11010001AZ

Imported by:
Zulat Pharmacy Sdn Bhd, 23 & 23A,
Jalan Bandar 3, Taman Melawati,
53100 Kuala Lumpur, Malaysia.

torrent

PHARMA

Manufactured by :
TORRENT PHARMACEUTICALS LTD.
Plot No. 810, Sector III, Industrial Area,
Pithampur, District Dhar - 454 775 (M.P.) INDIA

xxxxxxxx-5253

PRODUCT NAME	: Quace Cream	COUNTRY : Malaysia	LOCATION : Pithampur	Supersedes A/W No.: ZQCEXX5-MY			
ITEM / PACK	: Insert	NO. OF COLORS: 1	REMARK :				
DESIGN STYLE	: -	PANTONE SHADE NOS.:	SUBSTRATE :				
CODE	: xxxxxxxxx-5253		Activities	Department	Name	Signature	Date
DIMENSIONS (MM)	: 130 L x 180 W	Black C	Prepared By	Pkg.Dev			
ART WORK SIZE	: S/S		Reviewed By	Pkg.Dev			
DATE	: 21-01-2021		Approved By	Quality			

This colour proof is not colour binding. Follow Pantone shade reference for actual colour matching.

PRODUCT NAME	: Quace Cream	COUNTRY : Malaysia	LOCATION : Pithampur	Supersedes A/W No.: ZQCEXX5-MY			
ITEM / PACK	: Insert	NO. OF COLORS: 1	REMARK :				
DESIGN STYLE	: -	PANTONE SHADE NOS.:	SUBSTRATE :				
CODE	: xxxxxxxxx-5253		Activities	Department	Name	Signature	Date
DIMENSIONS (MM)	: 130 L x 180 W	Black C	Prepared By	Pkg.Dev			
ART WORK SIZE	: S/S		Reviewed By	Pkg.Dev			
DATE	: 21-01-2021		Approved By	Quality			

This colour proof is not colour binding. Follow Pantone shade reference for actual colour matching.