

CARNATION FOOTCARE CORN CAPS 40% MEDICATED PLASTERS

Self adhesive plaster

MAL Reg. No.: MAL11030026X

Each plaster contains salicylic acid BP 40% (36mg).

Description

The product consists of a 22 x 57mm piece of Nylon Weftstretch Zinc Oxide plaster onto which is centrally positioned a 19mm wide vinyl adhesive tape with the adhesive surface facing upwards.

An adhesive red coloured felt ring is central to the plaster and contains pink Carnation corn paste.

The adhesive surfaces and paste are covered by two pieces of embossed unplasticised PVC film, one overlapping the other.

Indications

Relieves and removes corns.

Action and mode of mechanisms of actions

Salicylic acid is a keratolytic and acts upon the intercellular lipid bilayer, allowing the keratinised cells to be debrided. The felt ring relieves the pressure on the corn which gives immediate relief from pain. The salicylic acid in the ointment is slowly released into the skin which it softens (keratolysis). The adhesive strapping holds the ring and the ointment in place for two days while the ointment is working. After a few days application, usually about six days, the softened corn can be easily scraped away. Often the centre of a hard corn can be removed whole.

Pharmacology

The formulation is designed to provide a gradual release of salicylic acid from the ointment to the restricted area of the corn. It is absorbed into the intercellular space in the outer epidermis.

Toxicology

As for all salicylates, absorption of large amounts can lead to salicylates poisoning. There is no danger from the small amount of salicylic acid in a corn cap (36mg). No more than 3 corns to be treated at any one time, therefore maximum exposure is 108mg salicylic acid, of which only a very small amount will be absorbed. Occasionally individuals who are sensitive to the rubber based adhesive suffer from a rash under the plaster.

Contraindications

Diabetes and impaired blood circulation of the feet unless used with a doctor's permission. Application to inflamed or broken skin. Hypersensitivity to salicylic acid or any of the ingredients in the product.

Caution has been suggested if caustic preparations are used in patients with significant peripheral neuropathy.

Side effects/adverse reactions

Mild pain has been experienced.

Occasional sensitivity to the ingredients has been reported. Local irritation or dermatitis may occur.

Discontinue use and remove plaster if excessive discomfort or irritation is experienced, or if sensitivity develops.

Precautions/warning

Carnation Corn Caps should not be used on healthy skin, by diabetics without their doctor's permission and should not be applied to skin that is cracked.

Cracks in the skin may go very deep and will let in the salicylic acid. Salicylic acid is safe if applied to intact skin but when applied to bare tissue causes pain and inflammation. Old people are prone to these cracks in the skin and their tissues are less able to withstand the irritant action of the salicylic acid.

People who have diabetes may suffer from poor circulation in the feet, which often causes the skin of the toes to become hard, dry and cracked. This condition is easily confused with a corn and applying a corn cap will make matters worse. Therefore, diabetics should always check with their doctor to make sure it is safe to

use a Carnation Corn Cap. Diabetics with poor circulation should have their feet regularly seen by a chiropodist.

Children under 15 years old and people with circulatory disorders should seek professional medical advice before use.

CONTAINS PEANUT OIL. DO NOT USE IF YOU ARE ALLERGIC TO PEANUT OR SOYA.
--

Symptoms and treatment for overdosage and antidote

None available.

Recommended dosage

Apply one corn plaster every two days, for ten days. A maximum of 5 plasters to be applied per corn. No more than 3 corns to be treated at any one time.

Instructions for use

First clean and thoroughly dry the skin around the corn. This is most important to ensure adhesion. Remove the facing material and position the felt ring very carefully so that it surrounds the corn. Firmly fix in position with the adhesive straps. When properly applied, it should not move or slip.

To obtain the best results, and in the interests of foot hygiene, change the corn cap every two days. Do not remove the corn until after the sixth day. If the corn is not easily removed, fit another cap and inspect every two days, changing the cap on each occasion.

Shelf life

Three years from the date of manufacture, as printed on the external wallet (package).

Packing

Each retail wallet contains 5 pieces of corn plaster. Store at less than 30°C, away from direct sources of heat and humidity.

Manufactured by :
Cuxson Gerrard & Co. Ltd.
125 Broadwell Road, Oldbury
West Midlands B69 4BF
Great Britain

Imported by :
Takaria (M) Sdn Bhd
Batu 4, Jalan Kelang Lama
PS-2A, Taman Evergreen
58100 Kuala Lumpur, Malaysia

Date of Revision: April 2014

R20099