

FEXOVID 180 mg TABLET

VIFEXfa-PC (MY)

DESCRIPTION

Oblong, peach film-coated tablet plain on both sides with bevel-edged and shallow convex faces.

COMPOSITION

Each film-coated tablet contains Fexofenadine Hydrochloride 180 mg.

PHARMACODYNAMICS

Fexofenadine is a non-sedating antihistamine with selective peripheral H₁-receptor antagonist activity and is a pharmacologically active metabolite of terfenadine. However, it does not possess significant sedative or antimuscarinic actions.

PHARMACOKINETICS

Absorption

Fexofenadine hydrochloride has rapid absorption following oral administration. The pharmacokinetics of fexofenadine are linear for oral doses up to 240 mg a day (120 mg twice a day).

Distribution

The volume of distribution is 5.4 to 5.8 liters/kilogram.

Protein binding

Fexofenadine has 60 to 70% plasma protein binding, predominantly to albumin and alpha₁-acid glycoprotein.

Biotransformation

About 5% of the total dose is metabolized; approximately 0.5 to 1.5% by cytochrome P450 3A4 isoenzyme metabolism and 3.5% transformed to a methyl ester metabolite by intestinal microflora.

Elimination

Elimination half-life of 14.4 hours in healthy subjects. The mean elimination half-life was 59% and 72% longer than healthy subjects for patients with mild renal impairment (creatinine clearance of 41 to 80 mL per minute) and severe renal impairment (creatinine clearance of 11 to 40 mL per minute) respectively. In patients on dialysis, half-life was 31% longer than in healthy subjects.

Excretion is mainly in the faeces with only 10% being present in the urine.

INDICATIONS

- Relief of symptoms associated with seasonal allergic rhinitis such as sneezing, rhinorrhea, itchy eyes, nose, and throat; and red, watery eyes.
- Treatment of uncomplicated skin manifestations of chronic idiopathic urticaria. It significantly reduces pruritus and the number of wheals.

CONTRAINDICATIONS

- Hypersensitivity to fexofenadine.
- Patients with renal function impairment.

PRECAUTIONS

Fexofenadine hydrochloride should be administered with care in the elderly and renally or hepatically impaired patients.

Patients with a history of or ongoing cardiovascular disease should be warned that, antihistamines have been associated with the adverse events, tachycardia and palpitations.

Carcinogenicity

Fexofenadine showed no carcinogenic potential in 18- and 24-month studies in mice and rats given oral terfenadine doses of 50 and 150 mg per kg of body weight (mg/kg) per day, respectively. These doses resulted in area under the plasma concentration-time curve (AUC) values for fexofenadine of up to four times the human therapeutic value based on the recommended dosage.

Mutagenicity

Fexofenadine was not mutagenic in *in vitro* bacterial or animal studies and *in vivo* animal studies.

PREGNANCY AND LACTATION

Adequate and well-controlled studies in humans have not been done.

It is not known whether fexofenadine is distributed into breast milk.

DRUG INTERACTIONS

Combinations containing any of the following medications, depending on the amount present, may also interact with this medication.

- Erythromycin or Ketoconazole.
- Antacids, aluminium and magnesium hydroxide-containing medication

MAIN SIDE/ADVERSE EFFECTS

Rare incidence: back pain, coughing, dizziness, drowsiness, dysmenorrhea, dyspepsia, fatigue, fever, headache, nausea, otitis media, sinusitis, viral infection (cold and flu).

OVERDOSE

Clinical features

Dizziness, drowsiness, dry mouth.

Treatment of overdosages

- There is no known antidote to fexofenadine.
- Treatment is generally symptomatic and supportive, may consider standard measure to remove any unabsorbed drug.
- Hemodialysis does not effectively remove fexofenadine from blood.
- Patients whom intentionally overdose should be referred for psychiatric evaluation.

DOSAGE AND ADMINISTRATION

Oral.

Adult and children aged 12 years and above:
One tablet once daily.

The efficacy and safety has not been studied in children under 12.

Storage: Store below 30°C.

Presentation/Packing:

Alu-alu blister pack of 4 x 7's.

Product Registration Holder: HOVID Bhd.
121, Jalan Tunku Abdul Rahman
(Jalan Kuala Kangsar),
30010 Ipoh, Perak, Malaysia.

Manufactured by: HOVID Bhd.
Lot 56442, 7 1/2 Miles, Jalan Ipoh/Chemor,
31200 Chemor, Perak, Malaysia.

Revision date: January 2018

hovid