

LIGNORAL Oral Spray 10% w/v

Lidocaine base
Multidose oral Spray
For surface anesthesia of oral mucous membrane

**For Medical
Professionals
Only**

Composition:

Each 100 ml of **LIGNORAL** Oral Spray contains:

Active ingredient:

Lidocaine 10 g

Excipients:

Saccharine sodium, Propylene glycol, Ethanol, Banana flavor, Purified water .

Product Description:

Amber glass bottle fitted in the top with spray-head and provided with oral applicator. The bottle contains clear colorless to pale yellow solution with sweet taste and banana flavor.

Properties:

LIGNORAL (lidocaine) is a safe and effective local anesthetic, belonging to the amide class of local anesthetics for surface anesthesia of oral mucous membrane.

Action:

Following local application of **LIGNORAL** oral spray, lidocaine penetrates oral mucous membrane altering cell membrane permeability of ions particularly sodium influx and preventing initiation and conduction of nerve impulses. Clinically, **LIGNORAL** oral spray provides prompt and profound local anesthesia that appears within 1 - 5 minutes post application to mucous membrane and lasts for about 15 - 30 minutes. This is beneficial in instantaneous relief of pain associated with dental practice or before otorhinolaryngology. Also, **LIGNORAL** action facilitates insertion of instruments during endotracheal intubation and endoscopic procedures of the airways and upper GIT. **LIGNORAL** oral spray is free of propellant gases and easy to handle besides high safety profile which recommends its use in all disciplines requiring oral mucous membrane surface anesthesia.

Pharmacokinetics:

Following application of **LIGNORAL** oral spray, absorption of lidocaine from mucous membrane depends upon concentration, total dose administered, the specific site of application and duration of exposure. About 64% of the absorbed lidocaine is bound to plasma proteins. Lidocaine is metabolized rapidly by the liver and excreted in the urine as metabolites and as unchanged drug.

Indications:

LIGNORAL oral spray is indicated in surface anesthesia:

- Before injections in dental practice, dental impressions (especially copperband impression), X-ray photography, removal of calculus.
- Before puncture of the maxillary sinus and surgical procedures in the nasal cavity, pharynx and epipharynx paracentesis.

Dosage and Mode of Administration:

Route of administration: Oral spray for topical use.

- In dental practice: 1 - 5 applications to oral mucous membranes. Onset of action is within 1 - 2 minutes.



0723171107

Item Name	Dimension (mm)	Colours #1	GTIN
Pamphlet Lignoral Topical Spray Export Malaysia	160 x 200 mm	black	
Dosage Form	Form	Varnish	Pharmacode
spray	Pamphlet	☐ Water base ☐ U.V ☐ S. U. V ☐ Solovan Matt ☐ V. Mutt	1153

Prepared by : Mahmoud Abd El Razek
Modified Date : 8.12.2025

- Puncture of the maxillary sinus and other otorhinolaryngology procedures: 3 applications. Onset of action is within 2 - 3 minutes.

Contraindications:

Hypersensitivity to lidocaine or any component of the oral spray. Lidocaine must not be used on open wounds or infected areas (local sepsis).

Warning and Precautions:

Local anesthetics must not be applied for prolonged periods or over large areas of mucous membranes or sprayed near or into the eyes. **LIGNORAL** oral spray must not be used by patients having allergy to local anesthetics and patients having epilepsy or liver, heart or kidney problems.

Side Effects:

Local adverse reactions might occur with applied lidocaine and include allergic reactions or drowsiness. Other side effects that may rarely occur include dizziness, blurred vision, tremors, nervousness, wheezing or difficulty in breathing, chest pain, severe rash or itching and increased sweating.

Pregnancy and Lactation:

No harmful effects have been reported on the use of local anesthetics during pregnancy and lactation. However, **LIGNORAL** oral spray should be used during pregnancy under strict medical supervision.

Drug Interactions:

- **LIGNORAL** should be used with caution in patients receiving class I antiarrhythmic drugs (such as tocainide and mexiletine) since the toxic effects are additive and potentially synergistic.
- When **LIGNORAL** is used concomitantly with other products containing local anesthetic agents, the amount absorbed from all formulations must be considered.

Interference with Laboratory Tests:

None reported with local anesthetics.

Overdosage and Treatment:

Symptoms of overdosage include drowsiness, confusion, nervousness, ringing in ears, blurred vision, feeling hot or cold, numbness, muscle twitches, slowed breathing or respiratory failure. Treatment is symptomatic and supportive.

Package: A Carton box containing an amber type III glass bottle of 15 ml solution capped with white polypropylene sprayer with (LDPE) plastic dip tube capped with transparent polypropylene cap & inox pump + Two 7-cm oral white polypropylene actuator + insert leaflet.

Storage conditions:

Store below 30°C.

Keep out of the reach of children.

Manufactured by:

EUROPEAN EGYPTIAN PHARM. IND:

Address: Km 25 Alexandria-Cairo Desert Road 21111, Alexandria, Egypt.

Product Registration Holder : Pharco Health Sdn Bhd

Address: Unit No. 643, PKNS BizPoint No. 67B, Tingkat 2, Jalan Plumbum P7/P

Seksyen 7, 40000 Shah Alam Selangor, Malaysia

Tel: +603 5566 8571 Fax: +603 5525 0040

Revision date: 8.12.2025



EUROPEAN EGYPTIAN PHARM. IND.

Manufactured by:

EUROPEAN EGYPTIAN PHARM. IND.

ALEXANDRIA - EGYPT