

PROMEZINE SYRUP 1mg/ml

Ingredient(s):

Each ml contains :

Promethazine Hydrochloride.....1mg

Preservative(s):

Methyl paraben.....1.8mg

Propyl paraben.....0.2mg

Pharmacology (Summary of Pharmacodynamic and Pharmacokinetic) :

Promethazine is a long acting antihistamine with a mild atropine-like anticholinergic effect and because of its marked effect on the central nervous system, is of great value as an antiemetic, hypnotic, tranquilizer, a potentiator of anaesthetic, hypnotics, sedatives and analgesics.

Absorption: Promethazine is well absorbed from the GI tract. The onset of sedative effects occurs within 20 minutes following oral administration. The duration of sedative effects varies but may range from 2-8 hours depending on dose and route of administration.

Distribution: Promethazine is widely distributed in body tissues. Compared with other organs, lower concentrations of the drug are found in the brain, but this concentration is higher than the plasma concentration. Promethazine readily crosses the placenta and it is not known if the drug is distributed into milk.

Elimination: Promethazine is metabolized by the liver. The drug is excreted slowly in urine and feces mainly as inactive promethazine sulfoxide and glucuronides.

Indication(s):

1. As symptomatic relief in perennial and seasonal rhinitis, pruritus, urticaria, dermatographism, angioedema, allergic conjunctivitis and amelioration of allergic reaction to blood or plasma.
2. As an adjunct in preoperative sedation in surgery and obstetrics.
3. As an antiemetic
4. For short term use:
 - a. Sedation and treatment of insomnia in adults
 - b. As a paediatric sedative.

Dosage & Administration:

Oral administration.

Not for use in children under age of 2 years.

As an antihistamine in allergy

Children 2-5 years	Either 5-15 mg as a single dose or 5 mg twice daily Maximum daily dose 15mg
Children 5-10 years	Either 10-25 mg as a single dose or 5-10 mg twice daily. Maximum daily dose 25mg
Children over 10 years and adults (including elderly)	Initially 10 mg twice daily Increasing to a maximum of 20 mg three times daily as required.

As an antiemetic

Children 2-5 years	5 mg to be taken the night before the journey To be repeated after 6-8 hours as required
Children 5-10 years	10 mg to be taken the night before the journey To be repeated after 6-8 hours as required
Children over 10 years and adults (including elderly)	25 mg to be taken the night before the journey To be repeated after 6-8 hours as required

Short term sedation

Children 2-5 years	15 or 20 mg as a single night time dose
Children 5-10 years	20 or 25mg as a single night time dose
Children over 10 years and adults (including elderly)	25 or 50mg as a single night time dose

Contraindication(s):

1. Promethazine should not be used in patients in coma or suffering from CNS depression of any cause.
2. This product should not be given to patients with a known hypersensitivity to promethazine or to any of the excipient.
3. Promethazine should be avoided in patients taking monoamine oxidase inhibitors up to 14 days previously.

Warning and Precautions:

1. Promethazine should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression.
2. To be used with caution and doctor's advice in children 2 to 6 years of age (when used for treatment of cough and cold).
3. This product may thicken or dry lung secretions and impair expectoration. It should be used with caution in patients with asthma, bronchitis or bronchiectasis.
4. This product should be used with care in patients with severe coronary artery disease, narrow angle glaucoma, epilepsy or hepatic and renal insufficiency.
5. Caution should be exercised in patients with bladder neck or pyloro-duodenal obstruction.
6. The use of promethazine in children and adolescents with signs and symptoms suggestive of Reye's Syndrome should be avoided.
7. Promethazine may mask the warning signs of ototoxicity caused by ototoxic drugs e.g. salicylates. It may also delay the early diagnosis of intestinal obstruction or raised intracranial pressure through the suppression of vomiting.
8. This product should not be used for longer than 7 days without seeking medical advice.
9. Patients should not drive or operate heavy machinery if they feel drowsy.

Interaction with Other Medicaments:

Promethazine will enhance the action of any anticholinergic agent, tricyclic antidepressant, sedative or hypnotic. Alcohol should be avoided during treatment. This product may interfere with immunological urine pregnancy tests to produce false-positive or false-negative results. Promethazine should be discontinued at least 72 hours before the start of skin tests as it may inhibit the cutaneous histamine response thus producing false-negative results.

Pregnancy and Lactation:

This product should not be used in pregnancy unless the physician considers it essential. The use of promethazine in the 2 weeks prior to delivery is not recommended in view of the risk of irritability and excitement in the neonate. Available evidence suggests that the amount excreted in milk is insignificant. However, there are risks of neonatal irritability and excitement.

Side Effects:

Side effects may be seen in a few patients: drowsiness, dizziness, restlessness, headaches, nightmares, tiredness, and disorientation. Anticholinergic side effects such as blurred vision, dry mouth and urinary retention occur occasionally. Infants are susceptible to the anti-cholinergic effects of promethazine, while other children may display paradoxical hyper-excitability. The elderly are particularly susceptible to the anti-cholinergic effects and confusion due to promethazine. Other side effects include urticaria, rash, pruritus, anorexia, gastric irritation, palpitations, hypotension, arrhythmias, extrapyramidal effects, muscle spasms and tic-like movements of the head and face. Anaphylaxis, jaundice and blood dyscrasias including haemolytic anemia rarely occur. Photosensitive skin reactions have been reported. Strong sunlight should be avoided during treatment.

Symptoms and Treatment of Overdose:

Symptoms of severe over dosage are variable. They are characterized in children by various combinations of excitation, ataxia, in-coordination, athetosis and hallucination, while adults may become drowsy and lapse into coma. Convulsions may occur in both adults and children: coma or excitement may precede their occurrence. Tachycardia may develop. Cardiorespiratory depression is uncommon. If the patient is seen soon enough after ingestion, it should be possible to induce vomiting with ipecacuanha despite the antiemetic effect of promethazine; alternatively, gastric lavage may be used. Treatment is otherwise supportive with attention to maintenance of adequate respiratory and circulatory status. Convulsions should be treated with diazepam or other suitable anticonvulsant.

Storage condition:

Keep in a tight container. Store below 30°C, protect from light and moisture.

Shelf Life:

2 Years from the date of manufacture.

Product Description and Packing (s):

Brown to dark brown color solution with cola flavour.
Plastic bottle of 60ml, 100ml and 120ml.

Manufacturer : **Y.S.P. INDUSTRIES (M) SDN. BHD.** (192593U)

Factory : Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian, Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan, Malaysia.
Tel: 03-89251215 (4 Lines) Fax: 03-89251298.



Y. S. P.

Office : No. 18, Jalan Wan Kadir, Taman Tun Dr. Ismail,
60000 Kuala Lumpur, Malaysia.
Tel: 03-77276390 (12 Lines) Fax: 0377276701
Toll Free: 1 800 88 3027