



Leaflet

Product Description

White, round, biconvex, chewable, mint flavoured tablet.

Content

Each tablet contains Simethicone 120 mg

Pharmacodynamics

Simethicone is a liquid dimethicone activated with finely divided silicon dioxide to enhance the defoaming properties of the silicon. Simethicone acts in the stomach and intestines by altering the surface tension of gas and mucus bubbles enabling them to coalesce. Coalescence of gas bubbles accelerates the passage of gas through belching or passing of flatus.

Pharmacokinetics

Simethicone is pharmacologically inert and is not absorbed from the gastrointestinal tract nor does it interfere with gastric secretion or absorption of nutrients. Simethicone is excreted unchanged in faeces.

Indication

Air-X 120 Tablet is indicated for the relief of the discomfort and pain caused by excessive gas in the stomach and intestines as in flatulence, dyspepsia, meteorism, peptic ulcer, spastic or irritable colon, diverticulitis and post-operative gaseous distention. Air-X 120 Tablet also can be used as an antifoaming agent prior to gastroscopy and endoscopic examination to enhance visualization and prior to radiography of bowel to reduce gas shadows.

Recommended Dose

Patients are required to chew the tablet thoroughly before swallowing them to enable the medicine work faster and more completely.

Adult: 1 tablet after meals and at bedtime as needed. It is recommended that the total dose should not exceed 480mg (4 tablets) in each 24 hour period.

The dosage is adjusted according to the age and symptoms of patients.

In preparation for radiography, gastroscopy and endoscopy, 80 - 240mg of simethicone tablet has been reported to be used alone or as adjunctive therapy in bowel preparation on the evening before and on the day of procedure to reduce intraluminal gas and to enhance the visibility and efficacy of the said procedures.

Route of Administration

Oral administration

Contraindications

Air-X 120 Tablet should not be taken by patients who have previous allergic reaction to simethicone or any of its excipients. It is also contraindicated in patients with known or suspected intestinal perforation and obstruction.

Warnings and Precautions

Air-X 120 Tablet contains glucose and sucrose, and as such, diabetic patients must be cautioned accordingly. For diabetics, Air-X SF tablet is the alternative.

Interactions with Other Medicaments

Bioavailability of phenytoin is reduced slightly when taken concurrently with simethicone. Hence, take Air-X 120 Tablet 2 hours before or after phenytoin.

Levothyroxine

- 1) Interaction Effect: decreased absorption of levothyroxine
- 2) Summary: Concomitant use of levothyroxine and simethicone may decrease levothyroxine efficacy by binding to and delaying or preventing levothyroxine absorption, potentially resulting in hypothyroidism. If concurrent use of levothyroxine and simethicone is required, administration of these agents should be separated by at least 4 hours (Prod Info SYNTHROID(R) oral tablets, 2008; Prod Info LEVOTHYROXINE SODIUM(R) oral tablet, 2007).
- 3) Severity: moderate
- 4) Onset: unspecified
- 5) Substantiation: theoretical
- 6) Clinical Management: Concomitant use of levothyroxine and simethicone may result in reduced levothyroxine efficacy by binding to and delaying or preventing absorption of levothyroxine, which may result in hypothyroidism. If concurrent use is necessary, separate administration of these agents by at least 4 hours (Prod Info SYNTHROID(R) oral tablets, 2008; Prod Info LEVOTHYROXINE SODIUM(R) oral tablet, 2007). Consider monitoring TSH levels and/or other measures of thyroid function when simethicone is initiated or discontinued during levothyroxine treatment.
- 7) Probable Mechanism: unknown

Pregnancy and Lactation

Simethicone is not absorbed into the body and it is not likely to cause problems to pregnant woman or breastfeeding mother. However, please inform the doctor if you are pregnant or breastfeeding

Side Effects

Simethicone is generally well tolerated when taken as directed. The following side effects have been reported very rarely: bloating, heartburn, constipation and diarrhoea.

Symptoms and Treatment of Overdose

As simethicone is not absorbed into the body there is no likelihood of an overdose.

Storage Condition

Store below 30°C in tight container
Protect from the heat and light
Do not consume after expiry date
Keep out of reach of children
Jauhi daripada kanak-kanak

Pack sizes

Box of 100, 200, 500, 2000 tablets

Shelf Life

Please refer to the outer box

Revision date: September 2010

Manufactured by:



76, Moo 10 Salaya, Bangpasi Road, Narapirom, Banglane, Nakompathorn.
73130 Thailand