

Patient Product Information

Supradyn® Film Coated Tablet

What is Supradyn and when is it used?

Supradyn contains as its active ingredients balance amounts of all the important vitamins (apart from vitamin K which is an exceptional case), essential minerals and trace elements. Vitamins are essential substances which are supplied to the body in food. Finally trace elements are vital substances which are required by the body in minute quantities but which are not always supplied in the daily food in adequate amounts. Supradyn is used as a dietary supplement. Supradyn is obtainable as film-coated tablets. The film-coated tablets are best swallowed whole with some liquid during meals.

How should Supradyn be taken?

1 film-coated tablet daily administered orally with sufficient fluid, unless the doctor prescribes a different dose for you.

What should be kept in mind when taking Supradyn?

If you are diabetic: the film-coated tablet has no caloric value, thus it can be administered to diabetic patients without problem.

When must Supradyn not be taken?

- Known hypersensitivity to any ingredient in the drug product
- Iron and/or copper metabolism disorders
- Existing hypervitaminosis A
- Existing hypervitaminosis D
- Hypercalcaemia
- Severe hypercalciuria
- Impaired renal function.

When should you be especially careful while taking Supradyn?

- The recommended dosage must not be exceeded. Very high doses of some ingredients, in particular vitamin A, vitamin D, iron and copper, can be harmful to health;
- Patients receiving other single vitamins or multivitamin preparations, any other medication or those under medical care must consult a healthcare professional before taking this product. In particular, patients receiving vitamin K antagonists and/or any other anti-coagulation treatment must consult a healthcare professional prior to use;
- Supradyn film coated tablets contain 3.5 mg sodium per tablet. To be taken into consideration by patients on a controlled sodium diet;

Does Supradyn have any interaction with medicines or food?

When used as recommended no specific interactions are expected. Manifold potential interactions are reported in the literature for the single ingredients.

It has been reported that vitamin E and K containing drugs should be used with caution in patients receiving anticoagulants or drugs that affect platelet aggregation.

Thus, patients receiving any other medication or those under medical care should consult a physician before taking this medicinal product;.

May Supradyn be taken during pregnancy or while breast-feeding?

Supradyn can be used during pregnancy and lactation; however, the recommended dosage must not be exceeded. As for every medicinal product, please consult your doctor.

Vitamin A doses of more than 10'000 IU per day have been found to be teratogenic if administered during the first trimester of pregnancy. Thus, Supradyn® must not be taken together with other drugs containing vitamin A, the synthetic isomers isotretinoin and etretinate, or beta-carotene, since large doses of the latter mentioned compounds are considered harmful to the fetus.

Permanent overdose of vitamin D might be harmful to the fetus and neonate. This must be taken into consideration if the infant is receiving any respective supplements.

What are the possible undesirable effects of Supradyn?

In general, this product is well-tolerated. However, in rare cases gastrointestinal symptoms (e.g. abdominal discomfort, constipation, vomiting, diarrhea, and nausea) might occur.

In very rare cases this product may cause allergic reaction. Symptoms may include hives, facial swelling, wheezing, skin reddening, rash, blisters, and shock. If an allergic reaction occurs, treatment must be stopped and a health care professional consulted.

You may occasionally notice a slight yellow discolouration of your urine. This effect is completely harmless and is due to the vitamin B2 contained in the preparation.

What are the possible signs of an overdose?

There is no evidence that this product can lead to an overdose when used as recommended.

Most, if not all reports concerning overdoses are associated with concomitant intake of high dosed single and/or multivitamin preparations. Acute or long-term overdose can cause hypervitaminosis A and D and hypercalcaemia as well as iron and copper toxicity.

Uncharacteristic initial symptoms, such as abrupt onset of headache, confusion, and gastrointestinal disturbances such as constipation, diarrhea, nausea, and vomiting might be indicative for an acute overdose.

If such symptoms occur, treatment must be stopped and a healthcare professional consulted.

How should Supradyn be stored?

The tablets should be stored in a dry place, and at a temperature below 25°C. The medicine should be kept out of the reach of children and should not be taken after the expiry date (EXP) shown on the pack. You can get further information from your doctor or pharmacist, who will have full prescribing information.

What does Supradyn contain?

12 vitamins	<i>Per film-coated tablet</i>
Vitamin A (1 mg)	3333 I.U.
Vitamin B1	4.5 mg
Vitamin B2	5.1 mg
Vitamin B6	6 mg
Vitamin B12	6 µg
Vitamin C	180 mg
Vitamin D3 (5 µg)	200 I.U.
Vitamin E	10 mg
Biotin (Vitamin H)	0.3 mg
Panthothenic Acid	21 mg
Folic Acid	0.2 mg
Nicotinamide	57 mg
8 minerals + trace elements	
Calcium	50 mg
Magnesium	40 mg
Phosphorus	50 mg
Chromium	10 µg
Copper	0.4 mg
Iron	3.6 mg
Manganese	0.5 mg
Zinc	3 mg

What packs are available?

Film-Coated Tablets

120s

Manufactured by

PT Bayer Indonesia
Jalan Raya Jakarta Bogor
Km 32 Cisalak, Kec.
Sukmajaya, Kota Depok,
Jawa Barat, Indonesia

Product Registration Holder

Bayer Co (Malaysia) Sdn. Bhd.
25-03 & 25-04, Level 25, IMAZIUM,
No. 8, Jalan SS21/37, Damansara
Uptown, 47400 Petaling Jaya,
Selangor

Date of revision: 01 June 2022



Made by P.T. Bayer Indonesia.
Manufactured under the authority of Bayer Consumer Care AG,
Basel, Switzerland.

BAYER